



7

**CLEAN SYNTHETIC VECTORS, PLASMIDS, TRANSGENIC PLANTS AND
PLANT PARTS CONTAINING THEM, AND METHODS FOR OBTAINING THEM.**

This application claims the priority of PCT Application No. PCT/IB00/01243, filed
September 4, 2000, and published in English on March 15, 2001, which application claims the
5 priority of French Patent Application No. 99111112, filed September 3, 1999.

The present invention relates to clean synthetic vectors, intended notably for use for
genetic transformation in the field of plant biotechnology.

In general, the vectors are known in the field of biotechnology and genetic manipulation.
The vectors which are currently used most commonly for genetic transformation, and in
10 particular in the field of plant biotechnology, present several disadvantages in their use, however.
In actual fact, and notably for plant transgenesis, use has often been made of a vector called
pBin19 (Frisch et al., 1995). The nucleotide sequence of this binary plasmid pBin19 is entirely
known. The problem, however, is that this plasmid is of large size (11.8 kbp) and that it contains
useless elements (more than half of pBin19) which are intolerable from a regulatory point of
15 view, or even detrimental to good replication. Moreover, the selection cassette of this plasmid is
located near the right border of the T-DNA. Thus, when the T-DNA is broken after the selection
cassette, it will not be possible to make any selection in order to retain solely the plants
possessing the expression cassette of the desired gene.

The expressions used in the description and the claims have the following meaning :

20 - "vector" means an expression system, for example DNA-coated projectiles, nucleic-
acid-based transit vehicles, nucleic acid molecules adapted to deliver nucleic acid, and
autonomous self-replicating circular DNA, for example plasmids, cosmids, phagemids, etc. If a
micro-organism or a recombinant cell culture is described as host of an "expression vector", this
can also include extrachromosomal circular DNA (such as for example mitochondrial or
25 chloroplast DNA), DNA which has been integrated into the host chromosome(s), where the
vector can be either replicated in a stable manner by the cells during mitosis as an autonomous
structure, integrated into the genome of the host, or maintained in the nucleus or cytoplasm of the
host.

- usually the vectors used for the genetic transformation exist in the form of plasmids. In this case, the "plasmid" is a molecule of autonomous circular DNA capable of replication in a cell. If a micro-organism or recombinant cell culture is described as the host of an "expression" plasmid, this comprises both extrachromosomal circular DNA molecules and DNA which has been integrated into the host chromosome(s). If the plasmid is maintained by a host cell, the plasmid is either replicated in a stable manner by the cells during mitosis as an autonomous structure, or integrated into the genome of the host;

- "clean" means that the vector comprises only sequences that are indispensable for its functionality and carries a nucleic acid sequence which comprises only elements that are indispensable for the expression of the host cell;

- "nucleic acid" means DNA or RNA ;

- "nucleic acid sequence" means a single- or double-stranded oligomer or polymer of nucleotide bases read from the 5' end towards the 3' end, and comprises self-replicating plasmids, genes, DNA or RNA polymers, which may or may not be infectious, and DNA, or RNA, either functional or non-functional. In the nucleotide notation used in the present application, unless specifically stated, the left end of a single-stranded nucleotide sequence is the 5' end;

- "derived nucleic acid sequence" means that the sequence derives directly or indirectly from the sequence referred to, for example by substitution, deletion, addition, mutation, fragmentation, and/or synthesis of one or more nucleotides;

- "promoter" means a nucleic acid region which is upstream of the translation initiation codon and which is involved in the recognition and binding of RNA polymerase and other transcription proteins;

- "plant promoter" is a promoter capable of initiating transcription in plant cells;

- "constitutive promoter" is a promoter capable of expressing nucleic acid sequences operationally bound to said promoter, in all or practically all the tissues of the host organism, during the whole development of said organism ;

- "tissue specific promoter" is a promoter capable of selectively expressing nucleic acid sequences operationally bound to said promoter, in certain specific tissues of the host organism;

- "operationally bound" means the binding of a functional or regulatory element, for example a promoter, to the nucleic acid sequence, or gene, to be expressed which codes for a protein to be produced, in such a way that this element influences the transcription of the bound nucleic acid sequence;

- "expression cassette" means nucleotide sequences capable of directing the expression of a nucleic acid sequence, or of a gene, coding for a polypeptide to be produced in a host organism compatible with such sequences. Such cassettes include at least one promoter and a transcription termination signal, and optionally other factors necessary or useful for the expression ;

- "heterologous sequence" or "heterologous nucleic acid sequence" means a sequence originating from a source, or from a species, foreign to the environment thereof, or if it originates from the same environment, which has been modified relative to its original form. The modification of the nucleic acid sequence can take place for example by treatment of the nucleic acid with a restriction enzyme in order to generate a nucleic acid fragment which can be operationally bound to a promoter. The modification can also take place by means of techniques such as directed mutagenesis;

- "box" means a nucleic acid sequence to which a regulatory function is attributed;

- "like" means that the box, and/or the nucleic acid sequence with which this term is associated, involves a certain sequence identity or consensus with a box and/or a known nucleic acid sequence, called a reference sequence, preferably a sequence identity of at least 50%, more preferably a sequence identity of at least 75%, and more particularly a sequence identity of at least 90% with the reference sequence. The percentage of sequence identity is calculated on the basis of a comparison window of at least 6 nucleotide bases. The determination of a comparison window can be carried out using sequence alignment algorithms to determine a homology with a reference sequence, for example the local homology algorithm, the homology alignment algorithm, and the similarity search algorithm, these algorithms existing also in computerized form, known by the names GAP, BESTFIT, FASTA and TFASTA. The percentage of sequence

identity is obtained by comparing the reference sequence with the box and/or the nucleic acid sequence;

- "situated" means the position on a nucleic acid sequence of an identified element, such as a "box", a restriction site, or a codon having a particular function. The position which is given by a figure refers to the position of the start of the element in the nucleic acid sequence, in the direction of reading of the latter, i.e. in the direction 5'→3';

- "transgenic plant" means a plant which has been obtained by genetic manipulation techniques, and covers the whole plants obtained, their progeny, and the plant organs, for example the roots, stems and leaves, obtained by these techniques. The transgenic plants according to the present invention can have different levels of ploidy, and can notably be polyploid, diploid, and haploid;

- "propagule" means a cluster or association of plant cells, which may or may not be structured, allowing the regeneration of a whole plant, for example explants, calli, stems, leaves, roots, cuttings, and even seeds.

The applicant of the present invention has succeeded, surprisingly, in producing clean synthetic vectors, in particular binary plasmids, of completely known nucleotide sequence, of small size, allowing the aforesaid disadvantages to be alleviated, and notably presenting a high replication rate relative to the existing vectors most commonly used. Furthermore, the applicant has succeeded at the same time in producing a range of vectors in such a way as to be able to choose the one which it is convenient to use according to the application envisaged and the environment of its use, and thus in such a way as to be able to better control the rate of expression of a gene to be expressed, coding for a polypeptide to be produced.

In addition, in some of the vectors according to the invention, each of the functional elements or components can be isolated by simple enzymatic digestion.

An object of the present invention is therefore a clean synthetic vector containing only the elements indispensable to its functionality and to the transgenesis of a cell, and notably of a plant cell.

According to a preferred embodiment, the vector comprises, as elements which are indispensable to its functionality and to the transgenesis of a cell, and which are operationally bound:

- at least one nucleic acid sequence coding for at least one first origin of replication, preferably an ori RK2, and more preferably at least one ori V of pRK2 of Escherichia coli with a broad host range;

- at least one nucleic acid sequence coding for a selection agent, preferably an antibiotic resistance gene, more preferably the npt III gene conferring resistance to kanamycin in bacteria;

- a trfA locus coding for at least one protein allowing an increase in the replication rate of the plasmid, preferably originating from pRK2, and more preferably coding for the proteins P285 and P382.

According to a more preferred embodiment the vector comprises the nucleic acid sequence identified by the number SEQ.ID01. Still more preferably, the vector consists of a single plasmid pMRT1105 whose nucleic acid sequence is identified by the number SEQ.ID01.

According to another preferred embodiment, the vector includes at least one nucleic acid sequence coding for a second origin of replication, preferably an ori of Escherichia coli, and more preferably an ori ColEI. More preferably, the vector comprises the nucleic acid sequence identified by the number SEQ.ID02, and still more preferably, the vector consists of a single plasmid pMRT1106 whose nucleic acid sequence is identified by the number SEQ.ID02.

Preferably, the synthetic vector according to the invention comprises a region (200 bp or less) made up of at least one nucleic acid sequence containing several unique enzymatic restriction sites called collectively «multiple cloning sites» (MCS).

According to a preferred embodiment, the vector also includes a nucleic acid sequence coding for a T-DNA comprising a right border RB and a left border LB, allowing the vector to act as a binary plasmid.

Preferably, the MCS is situated near (within 300 bp of) the right border, RB, of the T-DNA prior to the cloning of an insert into the MCS.

According to another preferred embodiment, the vector also includes a nucleic acid sequence coding for at least one expression promoter and at least one transcription terminator situated between the left border LB and the right border RB of the T-DNA. More preferably, the expression promoter is chosen from the group consisting of the constitutive promoters, the inducible promoters, the specific promoters, and preferably chosen from the plant expression promoters. Still more preferably, the expression promoter is chosen from the group consisting of the 35S CaMV promoter, the ep35S of CaMV, the pea plastocyanin gene promoter, its “enhancer” zones and derivatives, the wheat “High Molecular Weight Glutenin” (HMWG) promoter, the “Cassava Vein Mosaic Virus” CsVMV promoter, the “Commelina Yellow Mottle Virus” CoYMV promoter, the chimeric promoters of the CsVMV and CoYMV promoters, and their derivatives.

Preferably, the expression terminator is chosen from the functional terminators in a plant cell, and is preferably a 35S or nos terminator.

According to yet another preferred embodiment, the vector comprises at least one nucleic acid sequence coding for a selection agent that is functional in a plant cell, preferably at least one acid sequence coding for an antibiotic resistance gene, and/or a herbicide resistance gene.

Preferably, the sequence coding for a selection agent is a sequence coding for the bar («bialaphos resistance») resistance gene or pat («phosphinothricin acetyltransferase») gene, or else a sequence coding for the mutant or wild-type resistance gene nptII. More preferably still, the nucleic acid sequence coding for the selection agent is situated near the left border of the T-DNA.

According to another preferred embodiment of the present invention, the vector includes at least one expression cassette comprising an expression-promoting nucleic acid sequence operationally bound to a nucleic acid sequence to be expressed, coding for a polypeptide to be produced, itself bound to a transcription termination nucleic acid sequence. Preferably, the polypeptide to be produced is an enzyme or protein or derivative of the latter having activity in

vitro and/or in man and/or in animals, said activity comprising digestive, pancreatic, biliary, antiviral, anti-inflammatory, pulmonary, anti-microbial, nutritional, cosmetic, structural, blood, cardiovascular, ophthalmic, antigenic, immunostimulatory and cerebral activity. Examples of such proteins are for example the insulins, interferons, gastric, pancreatic or biliary lipases, elastases, antiproteases such as alpha-1 antitrypsin, structural proteins such as collagen, transferrins such as lactoferrin, proteins derived from blood, such as haemoglobin, human albumin and the blood cofactors, and antioxidants such as superoxide dismutase,

According to a particularly preferred embodiment of the invention, the vector is presented in the form of a binary, linear or circular plasmid, chosen from the group consisting of the nucleic acid sequences identified by the numbers SEQ.ID03, SEQ.ID04, SEQ.ID05, SEQ.ID06, SEQ.ID07, SEQ.ID08, SEQ.ID09, SEQ.ID10, SEQ.ID11, SEQ.ID12, SEQ.ID13, SEQ.ID14, SEQ.ID15, SEQ.ID16, SEQ.ID17, SEQ.ID18, SEQ.ID19, SEQ.ID20, SEQ.ID21 and SEQ.ID22.

According to a particularly advantageous embodiment, each functional component of the vector can be cleaved independently of the other components. Preferably, each functional component can be cleaved independently of the other components by enzymatic digestion at the level of a first unique restriction site and a second unique restriction site which are present in 1 vector.

Another object of the present invention is an isolated nucleic acid sequence, characterized in that it corresponds to a nucleic acid sequence chosen from the group consisting of the nucleic acid sequences identified by the numbers SEQ.ID01, SEQ.ID02, SEQ.ID03, SEQ.ID04, SEQ.ID05, SEQ.ID06, SEQ.ID07, SEQ.ID08, SEQ.ID09, SEQ.ID10, SEQ.ID11, SEQ.ID12, SEQ.ID13, SEQ.ID14, SEQ.ID15, SEQ.ID16, SEQ.ID17, SEQ.ID18, SEQ.ID19, SEQ.ID20, SEQ.ID21 and SEQ.ID22.

Yet another object of the present invention is a cell containing a vector or a nucleic acid sequence such as described earlier. The cell is preferably a plant cell.

Yet another object of the present invention is a transgenic plant having stably integrated in its genome a vector or a nucleic acid sequence such as described earlier. The plant is preferably chosen from the dicotyledon species, such as potato, tobacco, cotton, lettuce, tomato,

melon, cucumber, pea, rape, beetroot or sunflower, or the monocotyledon species, such as wheat, barley, oats, rice or maize.

Yet another object of the present invention is a propagule of a transgenic plant such as described earlier. The propagule is preferably a seed.

5 Yet another object of the present invention is a method for expression of a nucleic acid sequence, or gene, coding for a polypeptide to be produced, in a cell, characterized in that it comprises the stages consisting of:

- transforming the cell with a vector or a nucleic acid sequence such as described earlier;

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206

selection cassette near the left border, as well as rare restriction sites in their «polylinkers» (multiple cloning sites).

DESCRIPTION OF THE FIGURES

The invention will be understood better from the detailed description of the different embodiments which is given below in the form of examples, which are not limiting, and with reference to the drawing in the annex, in which:

- Figure 1 schematically represents a preferred minimal vector according to the present invention in the form of a plasmid identified by the reference pMRT1105, and with a length of 3508 base pairs;

- Figure 2 represents a schematic chart of a variant of the vector of Figure I in the form of a plasmid identified by the reference pMRT1106, and with a length of 4098 base pairs ;

- Figure 3 represents a schematic chart of a preferred vector according to the present invention, in the form of a binary plasmid identified by the reference pMRT1118, with a length of 5971 base pairs, and including a T-DNA comprising a selection cassette coding for resistance to an antibiotic, and a multiple cloning site (MCS);

- Figure 4 represents a preferred embodiment of the vector according to Figure 3, the multiple cloning site (MCS) including yet other unique sites, the vector being in the form of a binary plasmid identified by the reference pMRT1119, and with a length of 6016 base pairs ;

œ Figures 5 to 22 represent other preferred embodiments of the vector according to the invention, in the form of binary plasmids identified respectively by the references pMRT1121 (6017 base pairs), pMRT1122 (6016 base pairs), pMRT1155 (6017 base pairs), pMRT1175 (6767 base pairs), pMRT1176 (6767 base pairs), pMRT1191 (4805 base pairs), pMRT1192 (8654 base pairs), pMRT1193 (9143 base pairs), pMRT1195 (6865 base pairs), pMRT1196 (8654 base pairs), pMRT1201 (7943 base pairs), pMRT1202 (5614 base pairs), pMRT1203 (7503 base pairs), pMRT1204 (9390 base

pairs), pMRT1205 (7503 base pairs), pMRT1206 (9390 base pairs), pMRT1210 (10003 base pairs) and pMRT1212 (8987 base pairs) ;

5 ¶ Figure 23 graphically represents a comparison between the protein expression capacities of the sythetic vectors according to the present invention in comparison to a known vector system ;

¶ Figures 24 to 28 represent further embodiments of vectors according to the present invention, in the form of binary plasmids identified by the references pMRT1334 (9688 bp), pMRT1335 (15208 bp), pMRT1336 (9285 bp), pMRT1337 (8289 bp), pMRT1341 (14108 bp), pMRT1342 (15077 bp).

10 In the figures, the abbreviations have the following meanings :

20 - ori RK2 : ori V of pRK2 of Escherichia coli with a broad host range, unstable RK2 replicon producing a low number of copies;

30 - ori ColEI : the origin of replication ColEI of Escherichia coli;

40 - npt III gene: coding for neomycin phosphotransferase conferring resistance to kanamycin

50 - npt II gene: coding for neomycin phosphotransferase conferring resistance to kanamycin ;

60 - trfA : trfA locus (1481 bp) originating from pRK2 allowing the production of 2 proteins, P285 and P382, which enhance the replication of the plasmid by binding to the origin of replication ;

70 - Tnos : *nos* (nopaline synthetase) terminator;

80 - Pnos : *nos* (nopaline synthetase) promoter;

90 - LB : left border of a T-DNA ;

100 - RB : right border of a T-DNA ;

- MCS : multiple cloning site, S:XXX indicating the number of the starting base, E:XXX indicating the number of the end base of the MCS, the list starting with the first restriction site at the top and ending with the last restriction site at the bottom, in the direction 5'>3' ;

- polyA 35S : transcription termination (polyadenylation) signal ;

5 - gus gene: gene coding for beta-glucuronidase ;

- IA : the actin intron of rice ;

- PA : actin promoter of rice ;

- Phmwg : «high molecular weight glutenin» promoter of wheat ;

10 - bar («bialaphos resistance») gene: gene coding for the enzyme phosphinothricin acetyltransferase conferring resistance to glufosinate ;

- ep35S : «enhanced promoter» of the 35S ribosome.

EXAMPLES

15 In the detailed description which follows, all the enzymatic digestions by restriction enzymes were performed in accordance with the recommendations of the supplier New England Biolabs. The purifications with the aid of the «QIAquick Gel Extraction» and «QIAquick PCR Purification» kits were carried out in accordance with the recommendations of the supplier QIAGEN. The «Concert Rapid PCR Purification System» kit was used in accordance with the instructions of the supplier GIBCO BRL Life Technologies. The «GeneAmp PCR System 9700» thermocycler used is marketed by Perkin Elmer Applied Biosystems.

20 Example 1

1. Synthesis of pMRT1105

A clean synthetic vector according to the invention is presented preferably in the form of a minimal plasmid pMRT1105 (3508 bp) and is made up of the following elements:

- ori RK2 : ori V of pRK2 of Escherichia coli with a broad host range, unstable RK2 replicon producing a small number of copies (643 bp).

- npt III gene : confers resistance to kanamycin (bacterial selection marker, 1337 bp).

- TrfA : trfA locus (1481 bp) originating from pRK2, allowing the production of 2 proteins, P285 and P382, which enhance the replication of the plasmid by binding to the origin of replication.

The plasmid pMRT1105 results from the assembling of the fragments obtained by splicing overlap extension for «ori RK2» and a «part of npt III», of the fragment corresponding to a «part of trfA» produced by PCR («polymerase chain reaction») amplification and of «parts of trfA and npt III» isolated by enzymatic digestion.

1.1. Synthesis of the fragment carrying «ori RK2 and part of npt III»

The AvrII-StuI fragment (654 bp) carrying «ori RK2» (643 bp) was amplified by PCR from 5 ng of pBin19 matrix DNA with the aid of 20 pmoles of each of the 2 oligodeoxynucleotides, 5' AACCTAGGAAAAGACCGAGCGCCTTTGC 3' (SEQ.ID23) containing the AvrII restriction site and 5' CGGATTAATGGTAGAAGGCCTTTCACGGGAGGGTTCGAGAAGG 3' (SEQ.ID24) possessing the StuI restriction site, in the presence of 200 µM of each of the dNTPs, 60 mM Tris-SO₄ pH 9.1, 18 mM (NH₄)₂ SO₄, 1.8 mM MgSO₄ and 2 U of ELONGASE enzyme (GIBCO BRL Life Technologies) in a final reaction medium of 50 µl. The PCR amplification reaction was carried out in the «GeneAmp PCR System 9700» thermocycler. After denaturation at 94°C for 3 min., the DNA was subjected to 15 cycles each made up of the stages of denaturation at 94°C for 30 sec., hybridization at 55°C for 30 sec. and elongation at 68°C for 45 sec. Then, in the last cycle, the elongation was continued at 68°C for 3 min. Forty µl of the PCR reaction medium were then subjected to the action of 12.5 U of the Klenow fragment (New England Biolabs) in the presence of 2 µl of each of the dNTPs at 10 mM. The reaction was carried out at 37°C for 10 min. The PCR product so treated was then isolated by 2% agarose gel electrophoresis in TBE buffer (90 mM Tris-HCl, 2 mM Na₂-EDTA, 90 mM boric acid, pH 8.0) and purified with the aid of the «QIAquick Gel Extraction» kit. The DNA was recovered in 30 µl of H₂O.

The StuI-BstXI fragment (363 bp) carrying «part of npt III» (344 bp) was amplified and treated in the same way as the «ori RK2» fragment except that the 2 oligodeoxynucleotides used are 5' TGAAAGGCCTTCTACCATTAATCCGCGATAAACCCAGCGAACC 3' (SEQ.ID25) containing an StuI restriction site and 5'

5 ATGCATCCAAAATTTTGGTAGAATTTACAAGCTATAAGGTTATTGTCCTGGG 3' (SEQ.ID26) possessing the BstXI restriction site.

10 The 2 fragments carrying «ori RK2» and «part of npt III» were then assembled by splicing overlap extension in order to obtain the fragment «ori RK2 and part of npt III». To do this, a PCR amplification was performed from 7.5 µl of each of the treated PCR products corresponding to
AACCTAGGAAAAGACCGAGCGCCTTTGC 3' (SEQ.ID23) and 5'
ATGCATCCAAAATTTTGGTAGAATTTACAAGCTATAAGGTTATTGTCCTGGG 3' (SEQ.ID26)
in the presence of 200 µM of each of the dNTPs, 60 mM Tris-SO₄ pH 9.1, 18 mM (NH₄)₂ SO₄, 1.8 mM MgSO₄ and 2 U of ELONGASE enzyme in a final reaction medium of 50 µl. The PCR amplification reaction was carried out in the «GeneAmp PCR System 9700» thermocycler. After denaturation at 94°C for 3 min., the DNA was subjected to 15 cycles each made up of the stages of denaturation at 94°C for 30 sec., hybridization at 62°C for 30 sec. and elongation at 68°C for 45 sec. Then, in the last cycle, the elongation was continued at 68°C for 3 min. Forty µl of the PCR reaction medium were then subjected to 1.5% agarose gel electrophoresis in TBE buffer and
20 purified with the aid of the «QIAquick Gel Extraction» kit. The DNA, corresponding to the fragment carrying «ori RK2» and «part of npt III», was recovered in 30 µl of H₂O. Next, 27 µl of this DNA were hydrolysed by BstXI followed by AvrII, then purified with the aid of the «QIAquick PCR Purification» kit and recovered in 30 µl of H₂O.

1.2. Synthesis of the fragment carrying a «part of trfA»

25 The NdeI-AvrII fragment (295 bp) carrying «part of trfA» (295 bp) was amplified and treated in the same way as the fragment «ori RK2» except that the 2 oligodeoxynucleotides used are 5' ATCGACGAGGAAATCGTCGTGCTGTTTGC 3' (SEQ.ID27) situated upstream of the NdeI site and 5' AAACCTAGGAAATGCCAGTAAAGCGCTGGC 3' (SEQ.ID28) possessing the AvrII restriction site. The DNA fragments corresponding to a «part of trfA» originating from

the PCR amplification are then purified with the aid of the «QIAquick PCR Purification» kit, recovered in 30 µl of H₂O, hydrolysed by AvrII and NdeI, repurified with the aid of the «QIAquick PCR Purification» kit and recovered in 30 µl of H₂O.

1.3. Synthesis of the fragment carrying the «parts of trfA and npt III»

5 The BstXI-NdeI fragment (2196 bp) carrying the «parts of trfA and npt III» (2196 bp) was isolated from 9 µg of pBin19 DNA by enzymatic digestion by BstXI, purified with the aid of the «QIAquick PCR Purification» kit, then hydrolysed by NdeI. The DNA fragment was then isolated by 0.9% agarose gel electrophoresis in TBE buffer, purified with the aid of the «QIAquick Gel Extraction» kit and recovered in 30 µl of H₂O.

1.4. Obtaining pMRT1105

10 The 3 fragments, «ori RK2 and part of npt III», «parts of trfA and npt III» and «part of trfA», were ligated. To do this, the reaction mixture containing 7.5 µl of the digested fragments corresponding to «ori RK2 and part of npt III» and «parts of trfA and npt III», and 11 µl of the digested fragment corresponding to «part of trfA», was concentrated with the SpeedVac
15 AES1000 (SAVANT) in order to achieve a volume of 8 µl. After that, 1 µl of T4 DNA ligase x 10 buffer (New England Biolabs) and 400 units of T4 DNA ligase enzyme (New England Biolabs) were added. The ligation was carried out by a PCR reaction made up of 200 cycles each consisting of 2 stages, one at 30°C for 30 sec. and the other at 10°C for 30 sec., in the «GeneAmp PCR System 9700» thermocycler. The bacteria, Escherichia coli DH5⁺, which had previously
20 been made competent, were transformed (Hanahan, 1983). The plasmid DNA of the clones obtained, selected on Luria-Bertani medium (LB, bactotryptone 10 g/l, yeast extract 5 g/l, NaCl 10 g/l, Agar 15 g/l, pH 7.5) supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method (Birnboim and Doly, 1979) and verified by enzymatic digestions and by sequencing. The resulting plasmid selected was called pMRT1105 (3508 bp) and is represented
25 in Figure 1. Its complete sequence SEQ.ID01 is given in the sequence listing.

2. Synthesis of pMRT1106

The plasmid pMRT1106 (4098 bp) differs from pMRT1105 by the addition of the origin of replication ColE1 of *Escherichia coli* («ori ColE1»).

The fragment, carrying «ori ColE1» (590 bp), was isolated from the plasmid pBR322 marketed by New England Biolabs. The plasmid (5 µg) was digested by NdeI, purified with the aid of the «QIAquick PCR Purification» kit and recovered in 30 µl of TE buffer (10 mM Tris-HCl, 1 mM Na₂-EDTA, pH 8.0). The plasmid thus linearized was subjected to the action of 20 units of the Klenow fragment (New England Biolabs) in the presence of 12 µl of 500 mM Tris-HCl, pH 7.5 - 500 mM MgCl₂, 6 µl of 1M dithiothreitol, 6 µl of each of the 10 mM dNTPs in a reaction volume of 120 µl at 37°C for 30 min. The linearized plasmid so treated was purified with the aid of the «QIAquick PCR Purification» kit and recovered in 30 µl of H₂O. It was then digested by AlwNI in order to isolate the fragment carrying «ori ColE1», purified with the aid of the «QIAquick PCR Purification» kit and subjected to the action of 6 units of T4 DNA polymerase (New England Biolabs) in a reaction medium of 120 µl in the presence of 12 µl of T4 DNA polymerase x 10 buffer, 4 µl of 10 mM dNTP and 6 µl of BSA 1 mg/ml. The reaction was carried out at 37°C for 30 min. The DNA fragment was then isolated by 1.2% agarose gel electrophoresis in TBE buffer, purified with the aid of the «QIAquick Gel Extraction» kit and recovered in 50 µl of H₂O. This DNA fragment was then inserted into the plasmid pMRT1105 (2 µg) digested by StuI, purified with the aid of the «QIAquick PCR Purification» kit, dephosphorylated by 50 units of calf intestinal alkaline phosphatase (New England Biolabs) in a final reaction medium of 120 µl in the presence of 12 µl of 3 x 10 buffer (New England Biolabs) at 37°C for 1 hour, and purified with the aid of the «QIAquick PCR Purification» kit. The ligation by PCR reaction was carried out with 10 ng of digested dephosphorylated pMRT1105 plasmid and 100 ng of DNA fragments carrying «ori ColE1» in a reaction medium of 10 µl in the presence of 1 µl of T4 DNA ligase x 10 buffer (New England Biolabs) and 400 units of T4 DNA ligase enzyme (New England Biolabs). It is made up of 180 cycles consisting of 2 stages, one of 10°C for 30 sec. and the other of 30°C for 30 sec. in the «GeneAmp PCR System 9700» thermocycler. The bacteria, *Escherichia coli* DH5⁺, which had previously been made competent, were transformed (Hanahan, 1983). The validity of the clones obtained was verified by a PCR

test on bacterial colonies using 10 pmol of each of the 2 oligodeoxynucleotides, 5' AACCTAGGAAAAGACCGAGCGCCTTTGC 3' (SEQ.ID23) and 5' ATGCATCCAAAATTTTGGTAGAATTTACAAGCTATAAGGTTATTGTCCTGGG 3' (SEQ.ID26), in a reaction medium of 24 µl in the presence of 22 µl of SuperMix PCR (GIBCO BRL Life Technologies). The PCR amplification reaction was carried out in the «GeneAmp PCR System 9700» thermocycler. After denaturation at 94°C for 5 min., the DNA was subjected to 30 cycles each made up of the stages of denaturation at 94°C for 30 sec., hybridization at 60°C for 30 sec. and elongation at 72°C for 30 sec. Then, in the last cycle, the elongation was continued at 72°C for 7 min. The PCR reaction medium was then subjected to 1% agarose gel electrophoresis in TBE buffer. The insertion of the sequence «ori ColE1» was visualized by the presence of a fragment of about 1.6 kbp. The plasmid DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method (Birnboim and Doly, 1979) and verified by enzymatic digestions and by sequencing. The resultant plasmid selected was called pMRT1106 (4098 bp) and is represented in Figure 2. Its complete sequence SEQ.ID02 is given in the sequence listing.

3. Synthesis of pMRT1118

The plasmid pMRT1118 (5971 bp) differs from pMRT1106 by the introduction of a clean transfer DNA (T-DNA) in pMRT1106. This clean T-DNA is made up of the expression cassette of the mutant nptII gene (Frisch et al., 1995) under the control of the promoter and terminator of the nopaline synthase gene (nos, Depicker et al., 1982) of *Agrobacterium tumefaciens* placed between the right (RB) and left border (LB) of the plasmid pTiT37 of *Agrobacterium tumefaciens* nopaline strain.

3.1 Synthesis of the fragment carrying «end nptII - Tnos - LB»

3.1.1. Obtaining the fragment carrying LB

The AvrII-SmaI fragment (173 bp) carrying LB (151 bp) was amplified by PCR from 5 ng of pBin19 matrix DNA with the aid of 20 pmoles of each of the 2 oligodeoxynucleotides, 5' TTCCTAGGTTGACGTCTTCTGATGGGCTGCCTGTATCG 3' (SEQ.ID29) containing the AvrII and AatII restriction sites, and 5'

CCTATGGATATCCCCCGGGGATAGCCCCAGTACATTAAAAACGTCC 3' (SEQ.ID30) possessing the SmaI restriction site, in the presence of 200 µM of each of the dNTPs, 60 mM Tris-SO₄ pH 9.1, 18 mM (NH₄)₂ SO₄, 1.8 mM MgSO₄ and 2 U of ELONGASE enzyme (GIBCO BRL Life Technologies) in a final reaction medium of 50 µl. The PCR amplification reaction was carried out in the «GeneAmp PCR System 9700» thermocycler. After denaturation at 94°C for 3 min., the DNA was subjected to 15 cycles each made up of the stages of denaturation at 94°C for 30 sec., hybridization at 55°C for 30 sec. and elongation at 68°C for 45 sec. Then, in the last cycle, the elongation was continued at 68°C for 3 min. Forty µl of the PCR reaction medium were then subjected to the action of 12.5 U of the Klenow fragment (New England Biolabs) in the presence of 2 µl of each of the dNTPs at 10 mM. The reaction was carried out at 37°C for 10 min. The PCR product so treated was then isolated by 2% agarose gel electrophoresis in TBE buffer and purified with the aid of the «QIAquick Gel Extraction» kit. The DNA was recovered in 50 µl of H₂O.

3.1.2. Obtaining the fragment carrying the nos terminator (Tnos)

The SmaI-BspEI fragment (288 bp) carrying the nos terminator (256 bp) was amplified and treated in the same way as the fragment carrying LB except that the 2 oligodeoxynucleotides used are 5'CTATCCCCCGGGGATATCCATAGGCCCGATCTAGTAACATAGATGAC 3' (SEQ.ID31) containing a SmaI restriction site and 5' GCGCACTTGGGCCCATAGCTCGACGAACGATCGTTCAAACATTTGGC 3' (SEQ.ID32), possessing the Bsp120I restriction site.

3.1.3. Obtaining the fragment carrying «end of the nptII» gene («end nptII»)

The fragment (221 bp) carrying «end nptII» (221 bp) was amplified and treated in the same way as the fragment carrying LB except that the 2 oligodeoxynucleotides used are 5' TTCGTCGAGCTATGGGCCCAAGTGCGCATCCCGTGGGCGAAGAACTC 3' (SEQ.ID33) containing a Bsp120I restriction site and 5' TTCTTGACGAGTTCTTCTGAGCGGG 3' (SEQ.ID34) downstream of BstBI.

3.1.4. Obtaining the fragment «end nptII - Tnos - LB»

The fragments, «LB», «Tnos» and «end nptII», were assembled by splicing overlap extension in 2 stages :

- assembly of the fragments «LB» and Tnos» resulting in a fragment «Tnos - LB»,

5 - assembly of the fragment «Tnos - LB» and of the fragment «end nptII» resulting in a fragment «end nptII - Tnos -LB».

To do this, a first PCR amplification was carried out from 10 µl of each of the PCR products treated corresponding to «LB» and «Tnos», with the aid of 20 pmoles of each of the oligodeoxynucleotides, 5' TTCCTAGGTTGACGTCTTCTGATGGGCTGCCTGTATCG 3' (SEQ.ID29) and 5' GCGCACTTGGGCCCATAGCTCGACGAACGATCGTTCAAACATTTGGC 3' (SEQ.ID32), in the presence of 200 µM of each of the dNTPs, 60 mM Tris-SO₄ pH 9.1, 18 mM (NH₄)₂ SO₄, 1.8 mM MgSO₄ and 2 U of ELONGASE enzyme (GIBCO BRL Life Technologies) in a final reaction medium of 50 µl. The PCR amplification reaction was carried out in the «GeneAmp PCR System 9700» thermocycler. After denaturation at 94°C for 3 min., the DNA was subjected to 15 cycles each made up of the stages of denaturation at 94°C for 45 sec., hybridization at 62°C for 45 sec. and elongation at 68°C for 1 min. Then, in the last cycle, the elongation was continued at 68°C for 3 min. Forty µl of the PCR reaction medium were then subjected to the action of 12.5 U of the Klenow fragment (New England Biolabs) in the presence of 2 µl of each of the dNTPs at 10 mM. The reaction was carried out at 37°C for 10 min. The PCR product so treated, corresponding to the fragment «Tnos - LB» (477 bp), was then isolated by 2% agarose gel electrophoresis in TBE buffer and purified with the aid of the «QIAquick Gel Extraction» kit. The DNA was recovered in 30 µl of H₂O.

A second PCR amplification was then carried out from 7 µl of each of the PCR products treated, corresponding to «end nptII» and «Tnos - LB», with the aid of 20 pmoles of each of the oligodeoxynucleotides, 5' TTCCTAGGTTGACGTCTTCTGATGGGCTGCCTGTATCG 3' (SEQ.ID29) and 5' TTCTTGACGAGTTCTTCTGAGCGGG 3' (SEQ.ID34), in the presence of 200 µM of each of the dNTPs, 60 mM Tris-SO₄ pH 9.1, 18 mM (NH₄)₂ SO₄, 1.8 mM MgSO₄ and 2 U of ELONGASE enzyme (GIBCO BRL Life Technologies) in a final reaction medium of

50 µl. The PCR amplification reaction was carried out in the «GeneAmp PCR System 9700» thermocycler. After denaturation at 94°C for 3 min., the DNA was subjected to 15 cycles each made up of the stages of denaturation at 94°C for 45 sec., hybridization at 60°C for 45 sec. and elongation at 68°C for 1 min. Then, in the last cycle, the elongation was continued at 68°C for 3 min. This reaction is repeated 3 times for the PCR products treated, corresponding to «end nptII» and «Tnos - LB». The PCR reaction medium was then subjected to 2% agarose gel electrophoresis in TBE buffer and purified with the aid of the «QIAquick Gel Extraction» kit. The DNA fragment (672 bp), corresponding to «end nptII - Tnos - LB», was recovered in 100 µl of H₂O. Next, 95 µl of this DNA were hydrolysed by AvrII, purified with the aid of the «QIAquick PCR Purification» kit, digested by BstBI, purified with the aid of the «QIAquick PCR Purification» kit and recovered in 50 µl of H₂O. The 631 bp AvrII-BstBI fragment carries the sequence «end nptII – Tnos – LB».

3.2. Synthesis of the fragment carrying «end of the nos promoter (Pnos) and start of the nptII gene» («end Pnos - st. nptII»)

The fragment (1023 bp) carrying «end Pnos - st. nptII» was amplified and treated in the same way as the fragment carrying LB except that the 2 oligodeoxynucleotides used are 5' GGAATCGAAATCTCGTGATGGCAGG 3' (SEQ.ID39) and 5' ATTATTGCGCGTTCAAAAGTCGCC 3' (SEQ.ID40), and that the DNA was subjected to 15 cycles each made up of the stages of denaturation at 94°C for 45 sec., hybridization at 55°C for 45 sec. and elongation at 68°C for 1 min. The amplification was repeated for 2 samples of pBin19 DNA.

3.3. Synthesis of the fragment «end Pnos - nptII - Tnos - LB»

The fragments, «end Pnos - st. nptII» and «end nptII - Tnos - LB», were assembled by splicing overlap extension in order to produce a fragment «end Pnos - nptII - Tnos - LB».

To do this, a PCR amplification was carried out from 5 µl of each of the PCR products treated, corresponding to «end Pnos - st. nptII» and «end nptII - Tnos - LB», with the aid of 20 pmoles of each of the oligodeoxynucleotides, 5' TTCCTAGGTTGACGTCTTCTGATGGGCTGCCTGTATCG 3' (SEQ.ID29) and 5'

ATTATTGCGCGTTCAAAAGTCGCC 3' (SEQ.ID40), in the presence of 200 μ M of each of the dNTPs, 60 mM Tris-SO₄ pH 9.1, 18 mM (NH₄)₂ SO₄, 1.8 mM MgSO₄ and 2 U of eLONGASE enzyme (GIBCO BRL Life Technologies) in a final reaction medium of 50 μ l. The PCR amplification reaction was carried out in the «GeneAmp PCR System 9700» thermocycler. After
5 denaturation at 94°C for 3 min., the DNA was subjected to 15 cycles each made up of the stages of denaturation at 94°C for 45 sec., hybridization at 55°C for 45 sec. and elongation at 68°C for 1 min. Then, in the last cycle, the elongation was continued at 68°C for 3 min. This reaction was repeated 5 times for each of the PCR products treated, corresponding to «end Pnos - st. nptII» and «end nptII - Tnos - LB». The PCR reaction medium was then subjected to 0.7% agarose gel
10 electrophoresis in TBE buffer and purified with the aid of the «QIAquick Gel Extraction» kit. The DNA fragment (1608 bp) corresponding to «end Pnos - nptII - Tnos - LB», was recovered in 50 μ l of H₂O. Next, this DNA was subjected to the action of 62.5 U of the Klenow fragment (New England Biolabs) in the presence of 15 μ l of the Klenow fragment x 10 buffer (New England Biolabs) and 10 μ l of each of the dNTPs at 10 mM. The reaction was carried out at 37°C for 10 min. The DNA so treated was then purified with the aid of the «QIAquick PCR Purification» kit and recovered in 50 μ l of H₂O.

3.4. Synthesis of the fragment carrying «RB - MCS - st. Pnos»

3.4.1. Obtaining the fragment carrying «RB and multiple cloning site (MCS)» («RB - MCS»)

20 The fragment (210 bp) carrying «RB - MCS» was amplified and treated in the same way as the fragment carrying LB except that the 2 oligodeoxynucleotides used are 5' CGGTACCGAAGCTTTGAATTCACCTCGAGCAGATTGTCGTTTCCCGCC 3' (SEQ.ID35) possessing the restriction sites KpnI, HindIII, EcoRI and XhoI, and 5' TATCCTAGGAACCGGTAAACCCTGTGGTTGGCATGC 3' (SEQ.ID36) possessing the
25 restriction sites AvrII and AgeI.

3.4.2. Obtaining the fragment carrying «multiple cloning site (MCS) and start of the nos promoter (Pnos)» («MCS - st. Pnos»)

The fragment (209 bp) carrying «MCS - st. Pnos» was amplified and treated in the same way as the fragment carrying LB except that the 2 oligodeoxynucleotides used are 5'

5 ATATGAGACTCTAATTGGATACCGAGGGG 3' (SEQ.ID37) and 5' GCTCGAGTGAATTCAAAGCTTCGGTACCGTTGAAGGAGCCACTCAGCCG 3' (SEQ.ID38) possessing the restriction sites XhoI, EcoRI, HindIII and KpnI.

3.4.3. Obtaining the fragment carrying «RB - MCS - st. Pnos»

The fragments, «RB - MCS» and «MCS - st. Pnos», were assembled by splicing overlap extension in order to produce a fragment «RB - MCS - st. Pnos».

To do this, a PCR amplification was carried out from 12 µl of each of the PCR products treated, corresponding to «RB - MCS» and «MCS - st. Pnos», with the aid of 20 pmoles of each of the oligodeoxynucleotides, 5' ATATGAGACTCTAATTGGATACCGAGGGG 3' (SEQ.ID37) and 5' TATCCTAGGAACCGGTAAACCCTGTGGTTGGCATGC 3' (SEQ.ID36), in the presence of 200 µM of each of the dNTPs, 60 mM Tris-SO₄ pH9.1, 18 mM (NH₄)₂ SO₄, 1.8 mM MgSO₄ and 2 U of ELONGASE enzyme (GIBCO BRL Life Technologies) in a final reaction medium of 50 µl. The PCR amplification reaction was carried out in the «GeneAmp PCR System 9700» thermocycler. After denaturation at 94°C for 3 min., the DNA was subjected to 15 cycles each made up of the stages of denaturation at 94°C for 30 sec., hybridization at 60°C for 30 sec. and elongation at 68°C for 45 sec. Then, in the last cycle, the elongation was continued at 68°C for 3 min. This reaction was repeated for the remaining 3 x 12 µl of each of the PCR products treated, corresponding to «RB - MCS» and «MCS - st. Pnos». The PCR reaction medium was then subjected to 2% agarose gel electrophoresis in TBE buffer and purified with the aid of the «QIAquick Gel Extraction» kit (QIAGEN). The DNA fragment (390 bp) corresponding to «RB - MCS - st. Pnos», was recovered in 100 µl of H₂O. Next, 95 µl of this DNA was hydrolysed by AvrII, purified with the aid of the «QIAquick PCR Purification» kit, digested by Bsu36I, purified with the aid of the «QIAquick PCR Purification» kit and

recovered in 50 µl of H₂O. The 295 bp AvrII-Bsu36I fragment carries the sequence «RB – MCS – st. Pnos».

3.5. Synthesis of the T-DNA

The fragments, «RB - MCS - st. Pnos» and «end Pnos - nptII - Tnos - LB», were assembled by splicing overlap extension in order to produce a fragment «RB - MCS - Pnos - nptII - Tnos - LB» which corresponds to the T-DNA (1883 bp).

To do this, a PCR amplification was carried out from 4 µl of the PCR product treated corresponding to «RB - MCS - st. Pnos» and from 5 µl of the PCR product treated corresponding to «end Pnos - nptII - Tnos - LB», with the aid of 20 pmoles of each of the oligodeoxynucleotides, 5' TTCCTAGGTTGACGTCTTCTGATGGGCTGCCTGTATCG 3' (SEQ.ID29) and 5' TATCCTAGGAACCGGTAAACCCTGTGGTTGGCATGC 3' (SEQ.ID36), in the presence of 200 µM of each of the dNTPs, 60 mM Tris-SO₄ pH 9.1, 18 mM (NH₄)₂ SO₄, 1.8 mM MgSO₄ and 2 U of ELONGASE enzyme (GIBCO BRL Life Technologies) in a final reaction medium of 50 µl. The PCR amplification reaction was carried out in the «GeneAmp PCR System 9700» thermocycler. After denaturation at 94°C for 3 min., the DNA was subjected to 15 cycles each made up of the stages of denaturation at 94°C for 45 sec., hybridization at 55°C for 45 sec. and elongation at 68°C for 1 min. Then, in the last cycle, the elongation was continued at 68°C for 3 min. This reaction was repeated 10 times for each of the PCR products treated, corresponding to «RB - MCS - st. Pnos» and «end Pnos - nptII - Tnos - LB». The PCR reaction medium was then subjected to 0.8% agarose gel electrophoresis in TBE buffer and purified with the aid of the «QIAquick Gel Extraction» kit. The DNA fragment (1883 bp) corresponding to the T-DNA was recovered in 100 µl of H₂O. Next, this T-DNA was hydrolysed by AvrII (1873 bp fragment), purified with the aid of the «QIAquick PCR Purification» kit and recovered in 100 µl of H₂O.

3.6. Obtaining pMRT1118

The binary plasmid pMRT1118 (5971 bp) results from the introduction of the T-DNA fragment digested by AvrII into the AvrII site of the dephosphorylated pMRT1106 plasmid.

To do this, the pMRT1106 plasmid DNA (5 µg) was digested by AvrII, purified with the aid of the «QIAquick PCR Purification» kit, then dephosphorylated by 50 units of calf intestinal alkaline phosphatase (New England Biolabs) in a final reaction medium of 120 µl in the presence of 12 µl of 3 x 10 buffer (New England Biolabs) at 37°C for 1 hour, isolated by 0.6% agarose gel electrophoresis in TBE buffer, purified with the aid of the «QIAquick Gel Extraction» kit, dephosphorylated a second time with the calf intestinal alkaline phosphatase enzyme under the aforesaid conditions, and finally purified with the aid of the «QIAquick PCR Purification» kit and recovered in 50 µl of H₂O.

The ligation by PCR reaction was carried out with 32.5 ng of digested dephosphorylated pMRT1106 plasmid and 50 ng of digested T-DNA fragments in a reaction medium of 10 µl in the presence of 1 µl of T4 DNA ligase x 10 buffer (New England Biolabs) and 400 units of T4 DNA ligase enzyme (New England Biolabs). It is made up of 180 cycles consisting of 2 stages, one of 10°C for 30 sec. and the other of 30°C for 30 sec., in the «GeneAmp PCR System 9700» thermocycler.

The bacteria, *Escherichia coli* DH5⁺, which had previously been made competent, were transformed (Hanahan, 1983). The plasmid DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method (Birnboim and Doly, 1979) and verified by enzymatic digestions and by sequencing. The resultant plasmid selected was called pMRT1118 (Figure 3). Its complete sequence SEQ.ID03 is given in the sequence listing.

4. Synthesis of pMRT1119

The plasmid pMRT1119 (6016 bp) differs from pMRT1118 by the addition of additional unique restriction sites in the multiple cloning site MCS of pMRT1118.

4.1. Obtaining the additional unique restriction sites

The additional unique restriction sites (XbaI, SalI, PacI, BamHI, MluI, HpaI and FseI) were created by hybridization between the 2 oligodeoxynucleotides, 5'AGCTTGGCCGCGCGTTAACACGCGTGGATCCTTAATTAAGTCGACTCTAGAG 3' (SEQ.ID41) and 5'

AATTCTCTAGAGTCGACTTAATTAAGGATCCACGCGTGTTAACGGCCGGCCA 3' (SEQ.ID42). To do this, 5 µg of each of the 2 oligodeoxynucleotides were mixed and held at 85°C for 1 min., followed by a progressive reduction of the temperature to 80°C for 5 min., then a slow reduction of the temperature to 60°C in a water bath, and finally a rapid reduction of the temperature to the ambient temperature outside the water bath.

4.2. Obtaining pMRT1119

The binary plasmid pMRT1119 (6016 bp) results from the introduction of the sequence carrying the unique restriction sites into the HindIII and EcoRI sites of the pMRT1118 plasmid.

To do this, the pMRT1118 plasmid DNA (5 µg) was doubly digested by HindIII and EcoRI, purified with the aid of the «QIAquick PCR Purification» kit, and recovered in 50 µl of H₂O.

The ligation by PCR reaction was carried out with 75 ng of digested pMRT1118 plasmid and 500 ng of fragments carrying the unique restriction sites (described in 4.1.) in a reaction medium of 10 µl in the presence of 1 µl of T4 DNA ligase x 10 buffer (New England Biolabs) and 400 units of T4 DNA ligase enzyme (New England Biolabs). It is made up of 180 cycles consisting of 2 stages, one of 10°C for 30 sec. and the other of 30°C for 30 sec. in the «GeneAmp PCR System 9700» thermocycler.

The bacteria, *Escherichia coli* DH5⁺, which had previously been made competent, were transformed (Hanahan, 1983). The plasmid DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method (Birnboim and Doly, 1979) and verified by enzymatic digestions and by sequencing. The resultant plasmid selected was called pMRT1119 (Figure 4). Its complete sequence SEQ.ID04 is given in the sequence listing.

5. Synthesis of pMRT1121

The plasmid pMRT1121 (6017 bp) results from the insertion into pMRT1119 of a unique restriction site, BspEI, between the nopaline synthase promoter of *Agrobacterium tumefaciens* (Pnos) and the mutant nptII gene.

5 The BspEI site was inserted by the assembly by splicing overlap extension of two fragments obtained by PCR amplification.

5.1. Synthesis of the fragment carrying "part of nptII and BspEI site"

The 892 bp fragment carrying "part of nptII and BspEI site" was amplified by PCR from 5 ng of pBin19 matrix DNA with the aid of 20 pmoles of each of the 2 oligodeoxynucleotides, 5' GGAATCGAAATCTCGTGATGGCAGG 3' (SEQ.ID39) and 5' TAATCTGCATCCGGATCTGGATCGTTTCGC 3' (SEQ.ID43) carrying the BspEI site, in the presence of 200 µM of each of the dNTPs, 60 mM Tris-SO₄ pH 9.1, 18 mM (NH₄)₂ SO₄, 1.8 mM MgSO₄ and 2 U of eLONGase enzyme (GIBCO BRL Life Technologies) in a final reaction medium of 50 µl. This reaction was repeated. The PCR amplification reaction was carried out in a "GeneAmp PCR System 9700" thermocycler. After denaturation at 94°C for 3 min., the DNA was subjected to 15 cycles each made up of the stages of denaturation at 94°C for 45 sec., hybridization at 55°C for 45 sec. and elongation at 68°C for 1 min. Then, in the last cycle, the elongation was continued at 68°C for 3 min. The PCR product so obtained was isolated by 2% agarose gel electrophoresis in TBE buffer and purified with the aid of the «QIAquick Gel Extraction» kit. The DNA was recovered in 50 µl of H₂O.

5.2. Synthesis of the fragment carrying "BspEI site - MCS"

The fragment (250 bp) carrying "BspEI site - MCS" was amplified by PCR from 5 ng of pMRT1118 matrix DNA with the aid of 20 pmoles of each of the 2 oligodeoxynucleotides, 5' GCTCGAGTGAATTCAAAGCTTCGGTACCGTTGAAGGAGCCACTCAGCCG 3' (SEQ.ID38) and 5' ACGATCCAGATCCGGATGCAGATTATTTGG 3' (SEQ.ID44) carrying the BspEI site. The conditions of PCR amplification and treatment of the PCR fragment obtained are the same as described in 5.1.

5.3. Synthesis of the fragment carrying "part nptII - MCS"

The 1117 bp fragment carrying "part nptII - MCS" results from the assembly of the 2 PCR fragments, "part nptII - BspEI site" and "BspEI site - MCS", by splicing overlap extension.

To do this, a PCR amplification was carried out from 7.5 µl of each of the PCR products

5 treated, corresponding to "part nptII - BspEI site" and "BspEI site - MCS", with the aid of 20 pmoles of each of the 2 oligodeoxynucleotides, 5' GGAATCGAAATCTCGTGATGGCAGG 3' (SEQ.ID39) and 5'

GCTCGAGTGAATTCAAAGCTTCGGTACCGTTGAAGGAGCCACTCAGCCG 3'

(SEQ.ID38), in the presence of 200 µM of each of the dNTPs, 60 mM Tris-SO₄ pH 9.1, 18 mM

10 (NH₄)₂ SO₄, 1.8 mM MgSO₄ and 2 U of eLONGase enzyme (GIBCO BRL Life Technologies)

in a final reaction medium of 50 µl. This reaction was repeated 4 times. The PCR amplification

reaction was carried out in a "GeneAmp PCR System 9700" thermocycler. After denaturation at

94°C for 3 min., the DNA was subjected to 15 cycles each made up of the stages of denaturation

at 94°C for 45 sec., hybridization at 62°C for 45 sec. and elongation at 68°C for 1 min. Then, in

the last cycle, the elongation was continued at 68°C for 7 min. The PCR product so obtained was

isolated by 1% agarose gel electrophoresis in TBE buffer and purified with the aid of the

"QIAquick Gel Extraction" kit. The DNA was recovered in 50 µl of H₂O, digested by Bsu36I

and PstI, and subjected to 2% agarose gel electrophoresis in TEB buffer. The 315 bp DNA

fragment was isolated and purified with the aid of the "QIAquick Gel Extraction" kit.

5.4. Obtaining pMRT1121

The 315 bp DNA fragment Bsu36I - PstI, which carries the BspEI site, was ligated to the Bsu36I and PstI sites of the binary plasmid pMRT1119.

The plasmid pMRT1119 was previously digested by Bsu36I and PstI, subjected to 1% agarose gel electrophoresis in TEB buffer and purified with the aid of the «QIAquick Gel

25 Extraction» kit. The DNA was recovered in 50 µl of H₂O, dephosphorylated with 50 units of calf

intestinal alkaline phosphatase (New England Biolabs) in a final reaction medium of 120 µl in

the presence of 12 µl of 3 x 10 buffer (New England Biolabs) at 37°C for 1 hour, and purified

with the aid of the "QIAquick PCR Purification" kit.

The ligation by PCR reaction was carried out with 100 ng of digested dephosphorylated pMRT1119 plasmid and 30 ng of digested DNA fragments (315 bp) in a reaction medium of 10 µl in the presence of 1 µl of T4 DNA ligase x 10 buffer (New England Biolabs) and 400 units of T4 DNA ligase enzyme (New England Biolabs). It is made up of 180 cycles consisting of 2 stages, one of 10°C for 30 sec. and the other of 30°C for 30 sec., in a "GeneAmp PCR System 9700" thermocycler. The bacteria, *Escherichia coli* DH5 or SCS110 (deficient in Dam and Dcm methylases), which had previously been made competent, were transformed (Hanahan, 1983). The plasmid DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method and verified by enzymatic digestions and sequencing. The resultant plasmid was called pMRT1121 (Figure 5). Its complete sequence SEQ.ID05 is given in the sequence listing.

6. Synthesis of pMRT1122

The plasmid pMRT1122 (6016 bp) results from the introduction into pMRT1119 of a point mutation in the mutant *nptII* gene in order to restore the BglII restriction site and thus lead to the obtaining of the wild-type *nptII* gene.

To do this, the BglII restriction site was restored in the mutant *nptII* gene of pMRT1119 by directed mutagenesis in accordance with the Michael procedure (1994).

The oligodeoxynucleotide 5' ATGGGTCACGACGAGATCTTCGCCGTCGGG 3' (SEQ.ID45) was previously phosphorylated by subjecting 600 pmoles of the oligodeoxynucleotide to the action of 90 units of T4 kinase (Amersham) in the presence of 30 µl of T4 kinase x 10 buffer (Amersham) and 3 µl of 100 mM ATP in a final reaction medium of 300 µl at 37°C for 30 min. The oligodeoxynucleotide so treated was purified with the aid of the «Qiaquick Removal Nucleotide» (QIAGEN) kit in accordance with the supplier's recommendations.

The ligation by PCR was then carried out from 10 ng of pBIN19 matrix with the aid of 200 pmoles of each of the oligodeoxynucleotides, 5' GGAATCGAAATCTCGTGATGGCAGG 3' (SEQ.ID39), 5' ATGGGTCACGACGAGATCTTCGCCGTCGGG 3' (SEQ.ID45), which was phosphorylated, including the BglII restriction site, and 5'

ATTATTGCGCGTTCAAAAGTCGCC 3' (SEQ.ID40), in the presence of 400 μ M of each of the dNTPs, 10 μ l of Taq DNA ligase x 10 buffer (New England BioLabs), 5 units of Vent DNA Polymerase (New England BioLabs) and 40 units of Taq DNA ligase (New England BioLabs) in a final reaction medium of 100 μ l. The PCR ligation reaction was carried out in the «GeneAmp PCR System 9700» thermocycler by carrying out the following three successive phases: the first phase consisted of a cycle made up of the stages of denaturation at 94°C for 5 min., hybridization at 50°C for 1 min. and elongation at 65°C for 4 min.; the second phase was made up of 28 cycles each comprising the stages of denaturation at 94°C for 30 sec., hybridization at 50°C for 1 min. and elongation at 65°C for 4 min. ; and, finally, the last phase consisted of a cycle made up of the stages of denaturation at 94°C for 30 sec., hybridization at 50°C for 1 min. and elongation at 65°C for 15 min. The PCR reaction medium was then subjected to 0.8% agarose gel electrophoresis in TBE buffer and purified with the aid of the «QIAquick Gel Extraction» kit. The DNA fragments (1023 bp) so treated were hydrolysed by NcoI and PstI and subjected to 2% agarose gel electrophoresis. The 383 bp DNA fragments were isolated, purified with the aid of the «QIAquick Gel Extraction» kit, recovered in 50 μ l of H₂O and ligated to the NcoI and PstI sites of the pMRT1119 plasmid. To do this, the pMRT1119 plasmid DNA was digested by NcoI and PstI, purified on 0.8% agarose gel. The fragment corresponding to the plasmid was isolated , purified with the aid of the «QIAquick Gel Extraction» kit, then dephosphorylated with 50 units of calf intestinal alkaline phosphatase (New England Biolabs) in a final reaction medium of 120 μ l in the presence of 12 μ l of 3 x 10 buffer (New England Biolabs) at 37°C for 1 hour, and, finally, recovered in 50 μ l of H₂O.

The ligation by PCR was carried out with 50 ng of digested dephosphorylated pMRT1119 plasmid and all the DNA fragments digested and treated in a reaction medium of 10 μ l in the presence of 1 μ l of T4 DNA ligase x 10 buffer (New England Biolabs) and 400 units of T4 DNA ligase enzyme (New England Biolabs). It is made up of 180 cycles consisting of 2 stages, one of 10°C for 30 sec. and the other of 30°C for 30 sec., in the «GeneAmp PCR System 9700» thermocycler.

The bacteria, *Escherichia coli* DH5⁻, which had previously been made competent, were transformed (Hanahan, 1983). The plasmid DNA of the clones obtained, selected on LB medium

supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method (Birnboim and Doly, 1979) and screened by digestion by BglIII. The plasmid DNA of the clone selected was verified by enzymatic digestions and by sequencing. The resultant plasmid selected was called pMRT1122 (6016 bp). It is represented in Figure 6 and its complete sequence SEQ.ID06 is given in the sequence listing.

7. Synthesis of pMRT1155

The plasmid pMRT1155 (6017 bp) differs from pMRT1122 by the presence of the BspEI site.

To do this, the 315 bp insert fragment containing the BspEI site was obtained by digestion of pMRT1121 by Bsu36I and PstI, 1.2% agarose gel electrophoresis in TEB buffer, purification with the aid of the "QIAquick Gel Extraction" kit and recovery in 50 µl of H₂O.

The pMRT1122 vector fragment was obtained by digestion of pMRT1122 by Bsu36I and PstI, 1.2% agarose gel electrophoresis in TEB buffer, purification with the aid of the "QIAquick Gel Extraction" kit and recovery in 50 µl of H₂O. After that, digested pMRT1122 was dephosphorylated with 50 units of calf intestinal alkaline phosphatase (New England Biolabs) in a final reaction medium of 120 µl in the presence of 12 µl of 3 x 10 buffer (New England Biolabs) at 37°C for 1 hour, and purified with the aid of the "QIAquick PCR Purification" kit.

The ligation by PCR reaction was carried out with 100 ng of dephosphorylated digested pMRT1122 plasmid and 50 ng of digested DNA fragments (315 bp) in a reaction medium of 10 µl in the presence of 1 µl of T4 DNA ligase x 10 buffer (New England Biolabs) and 400 units of T4 DNA ligase enzyme (New England Biolabs). It is made up of 180 cycles consisting of 2 stages, one of 10°C for 30 sec. and the other of 30°C for 30 sec. in a "GeneAmp PCR System 9700" thermocycler. The bacteria, *Escherichia coli* DH5 or SCS110 (deficient in Dam and Dcm methylases), which had previously been made competent, were transformed (Hanahan, 1983). The plasmid DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method and verified by enzymatic digestions and sequencing. The resultant plasmid was called pMRT1155 (Figure 7). Its complete sequence SEQ.ID07 is given in the sequence listing.

8. Obtaining basic binary plasmids comprising the sequence "eP35S - T35S"

8.1. Obtaining pMRT1205

The plasmid pMRT1205 (7503 bp) includes the expression cassette of the mutant nptII gene and a "double 35S promoter - 35S terminator" sequence (eP35S - T35S) for the cloning of genes of interest. It results from the cloning of the eP35S promoter of the cauliflower mosaic virus (CaMV) into pMRT1175. The CaMV eP35S promoter corresponds to a duplication of the transcription-activating sequences situated upstream of the TATA element of the 35S promoter (Kay et al., 1987). As for the plasmid pMRT1175 (6767 bp), this results from the cloning of T35S of the cauliflower mosaic virus into pMRT1121. The transcription-terminating sequence, CaMV T35S, corresponds to the non-coding region in 3' of the circular double-stranded DNA sequence of the cauliflower mosaic virus producing the 35S transcript (Franck et al., 1980).

8.1.1. Obtaining pMRT1175

The plasmid pMRT1175 results from the cloning of the EcoRI - XhoI insert DNA fragments corresponding to the T35S into the EcoRI and XhoI sites of the vector pMRT1121 produced from Escherichia coli strain SCS110.

The pMRT1121 vector fragment was obtained by digestion of 7 µg of pMRT1121 by EcoRI and XhoI, purification with the aid of the "QIAquick PCR Purification" kit and recovery in 50 µl of H₂O. After that, digested pMRT1121 was dephosphorylated with 50 units of calf intestinal alkaline phosphatase (New England Biolabs) in a final reaction medium of 120 µl in the presence of 12 µl of 3 x 10 buffer (New England Biolabs) at 37°C for 1 hour, and purified with the aid of the "QIAquick PCR Purification" kit and recovered in 50 µl of H₂O.

The EcoRI – XhoI insert DNA fragments (757 bp) corresponding to the T35S (750 bp) were obtained by digestion by EcoRI and XhoI of the plasmid pJIT163, which derives from the plasmid pJIT60 (Guerineau and Mullineaux, 1993). They were then subjected to 1% agarose gel electrophoresis in TEB buffer, purified with the aid of the «QIAquick Gel Extraction» kit and recovered in 30 µl of H₂O.

The ligation by PCR reaction was carried out as described earlier in 7, with 80 ng of dephosphorylated digested pMRT1121 plasmid and 80 ng of digested insert DNA fragments. The bacteria, *Escherichia coli* DH5⁺, which had previously been made competent, were transformed. The plasmid DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method and verified by enzymatic digestions and sequencing. The resultant plasmid was called pMRT1175 (6767 bp). It is represented in Figure 8 and its complete sequence SEQ.ID08 is given in the sequence listing.

8.1.2. Obtaining pMRT1205

The plasmid pMRT1205 results from the cloning of the KpnI - HindIII insert DNA fragments corresponding to eP35S into the KpnI and HindIII sites of the vector pMRT1175.

The pMRT1175 vector fragment was obtained by digestion of 4.1 µg of pMRT1175 by KpnI and HindIII, purification with the aid of the «Concert Rapid PCR Purification System» kit and recovery in 100 µl of H₂O. After that, digested pMRT1175 was dephosphorylated with 50 units of calf intestinal alkaline phosphatase (New England Biolabs) in a final reaction medium of 120 µl in the presence of 12 µl of 3 x 10 buffer (New England Biolabs) at 37°C for 1 hour, purified with the aid of the «Concert Rapid PCR Purification System» kit, and recovered in 50 µl of H₂O.

The KpnI - HindIII insert DNA fragments (743 bp) corresponding to eP35S (735 bp) were obtained by digestion by KpnI and HindIII of the plasmid pJIT163, which derives from the plasmid pJIT60 (Guerineau and Mullineaux, 1993). They were then subjected to 1% agarose gel electrophoresis in TEB buffer, purified with the aid of the "QIAquick Gel Extraction" kit and recovered in 30 µl of H₂O.

The ligation by PCR reaction was carried out as described earlier in 7, with 48 ng of dephosphorylated digested pMRT1175 plasmid and 30 ng of digested insert DNA fragments. The bacteria, *Escherichia coli* DH5⁺, which had previously been made competent, were transformed. The plasmid DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method and verified by enzymatic digestions and

sequencing. The resultant plasmid was called pMRT1205 (7503 bp). It is represented in Figure 19 and its complete sequence SEQ.ID19 is given in the sequence listing.

8.2. Obtaining pMRT1203

The plasmid pMRT1203 (7503 bp) includes the expression cassette of the wild-type nptII gene and a "double 35S promoter - 35S terminator" sequence (eP35S - T35S) for the cloning of genes of interest. It results from the cloning of the eP35S promoter of the cauliflower mosaic virus into pMRT1176. As for the plasmid pMRT1176 (6767 bp), this results from the cloning of T35S of the cauliflower mosaic virus into pMRT1155. The eP35S promoter and the T35S terminator are described in 8.1.

8.2.1. Obtaining pMRT1176

The plasmid pMRT1176 results from the cloning of the EcoRI - XhoI insert DNA fragments (757 bp) corresponding to the T35S (750 bp) into the EcoRI and XhoI sites of the vector pMRT1155 produced from Escherichia coli strain SCS110.

The plasmid pMRT1176 was obtained in accordance with the methodologies described in 8.1.1., except that 6.3 µg of the vector pMRT1155, which constitutes the cloning vector, were digested and treated. The plasmid pMRT1176 (6767 bp) is represented in Figure 9. Its complete sequence SEQ.ID09 is given in the sequence listing.

8.2.2. Obtaining pMRT1203

The plasmid pMRT1203 results from the cloning of the KpnI - HindIII insert DNA fragments (743 bp) corresponding to eP35S (735 bp) into the KpnI and HindIII sites of the vector pMRT1176.

The plasmid pMRT1203 was obtained in accordance with the methodologies described in 8.1.2., except that 3.9 µg of the vector pMRT1176, which constitutes the cloning vector, were digested and treated. The plasmid pMRT1203 (7503 bp) is represented in Figure 17. Its complete sequence SEQ.ID17 is given in the sequence listing.

9. Obtaining binary plasmids comprising the sequence "eP35S - uidA - T35S"

These vectors were employed in transformation of tobacco in order to evaluate them and determine the incidence of the mutation of the nptII gene.

9.1. Obtaining pMRT1206

5 The plasmid pMRT1206 (9390 bp) includes the expression cassette of the mutant nptII gene and the expression cassette of the uidA gene (gus). It results from the cloning of the eP35S promoter of the cauliflower mosaic virus into pMRT1196. As for the plasmid pMRT1196 (8654 bp), this results from the cloning of the uidA gene (Jefferson RA et al., 1986) into pMRT1175.

9.1.1. Obtaining pMRT1196

The plasmid pMRT1196 results from the cloning of the insert DNA fragments (SmaI - "SacI + T4 DNA polymerase"), corresponding to the uidA gene, into the "XbaI + Klenow" site of the vector pMRT1175.

The pMRT1175 vector fragment was obtained by digestion of 10 µg of pMRT1175 by XbaI, purified with the aid of the "QIAquick PCR Purification" kit, recovered in 50 µl of H₂O and subjected to the action of 20 units of the Klenow fragment (New England Biolabs) in the presence of 12 µl of 500 mM Tris-HCl pH 7.5, 500 mM MgCl₂, 6 µl of 1M dithiothreitol, 6 µl of each of the 10 mM dNTPs in a reaction volume of 120 µl at 37°C for 30 min. It was then purified with the aid of the "QIAquick PCR Purification" kit and recovered in 50 µl of H₂O. The plasmid pMRT1175 so digested and treated was dephosphorylated with 50 units of calf intestinal alkaline phosphatase (New England Biolabs) in a final reaction medium of 120 µl in the presence of 12 µl of 3 x 10 buffer (New England Biolabs) at 37°C for 1 hour, and purified with the aid of the "QIAquick PCR Purification" kit and recovered in 50 µl of H₂O.

The insert DNA fragments (2 µg) corresponding to the uidA gene (1.8 kbp) were obtained by digestion of pBI221 (marketed by Clontech) by SacI, purification with the aid of the "QIAquick PCR Purification" kit and recovery in 50 µl of H₂O. After that, digested pBI221 was subjected to the action of 6 units of T4 DNA polymerase (New England Biolabs) in a reaction

medium of 120 µl in the presence of 12 µl of T4 DNA polymerase x 10 buffer, 4 µl of 10 mM dNTPs and 6 µl of BSA 1 mg/ml. The reaction was carried out at 37°C for 30 min. The plasmid pBI221 so treated was purified with the aid of the "QIAquick PCR Purification" kit and recovered in 50 µl of H₂O. Finally, pBI221 so treated was digested by SmaI. The «[SacI + T4 DNA polymerase] – SmaI» fragment (1882 bp) was isolated by 0.8% agarose gel electrophoresis, purified with the aid of the "QIAquick Gel Extraction" kit and recovered in 50 µl of H₂O.

The ligation by PCR reaction was carried out in accordance with the methodologies described in 7, with 15 ng of dephosphorylated digested pMRT1175 plasmid and 100 ng of digested insert DNA fragments. The bacteria, *Escherichia coli* DH5⁺, which had previously been made competent, were transformed. The plasmid DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method and verified by enzymatic digestions and sequencing. The resultant plasmid was called pMRT1196 (8654 bp). It is represented in Figure 14 and its complete sequence SEQ.ID14 is given in the sequence listing.

9.1.2. Obtaining pMRT1206

The plasmid pMRT1206 (9390 bp) results from the cloning of the KpnI – HindIII insert DNA fragments (743 bp), corresponding to eP35S (735 bp), into the KpnI and HindIII sites of the vector pMRT1196.

The plasmid pMRT1206 was obtained in accordance with the methodologies described in 8.1.2., except that 2 µg of the vector pMRT1196, which constitutes the cloning vector, were digested and treated. It is represented in Figure 24 and its complete sequence SEQ.ID24 is given in the sequence listing.

The plasmid pMRT1206 was introduced into *Agrobacterium tumefaciens* strain LBA4404 by direct transformation in accordance with the procedure of Holsters et al. (1978).

9.2. Obtaining pMRT1204

The plasmid pMRT1204 (9390 bp) includes the expression cassette of the wild-type nptII gene and the expression cassette of the uidA gene. It results from the cloning of the eP35S promoter of the cauliflower mosaic virus into pMRT1192. As for the plasmid pMRT1192 (8654 bp), this results from the cloning of the uidA gene into pMRT1176.

9.2.1. Obtaining pMRT1192

The plasmid pMRT1192 results from the cloning of the insert DNA fragments (SmaI - "SacI + T4 DNA polymerase"), corresponding to the uidA gene, into the "XbaI + Klenow" site of the vector pMRT1176.

The plasmid pMRT1192 (8654 bp) was obtained in accordance with the methodologies described in 9.1.2., except that the cloning vector is pMRT1176. It is represented in Figure 11 and its sequence SEQ.ID11 is given in the sequence listing.

9.2.2. Obtaining pMRT1204

The plasmid pMRT1204 (9390 bp) results from the cloning of the KpnI – HindIII insert DNA fragments (743 bp), corresponding to eP35S (735 bp), into the KpnI and HindIII sites of the vector pMRT1192.

The plasmid pMRT1204 was obtained in accordance with the methodologies described in 8.1.2., except that 2 µg of the vector pMRT1192, which constitutes the cloning vector, were digested and treated. It is represented in Figure 18 and its complete sequence SEQ.ID18 is given in the sequence listing.

The plasmid pMRT1204 was introduced into *Agrobacterium tumefaciens* strain LBA4404 by direct transformation in accordance with the procedure of Holsters et al. (1978).

10. Obtaining binary plasmids containing the bar gene

The final plasmids were employed in transformation of maize and/or tobacco in order to evaluate them.

10.1. Creation of pMRT1210

The plasmid pMRT1210 (10003 bp) includes the expression cassette of the bar gene and the expression cassette of the uidA gene. It results from the cloning of the expression cassette of the uidA gene into pMRT1195. As for the plasmid pMRT1195 (6865 bp), this results from the cloning of the sequence "promoter followed by intron 1 of the rice actin gene" (McElroy D et al., 1991) into pMRT1191. The plasmid pMRT1191 (4805 bp) derives from pMRT1119.

10.1.1. Obtaining pMRT1191

The plasmid pMRT1191 (4805 bp) differs from pMRT1119 by the deletion of the nos promoter and mutant nptII gene sequences.

It was obtained by digestion by Bsp120I of 10 µg of pMRT1119, followed by purification with the aid of the "QIAquick PCR Purification" kit and recovery in 50 µl of H₂O. After that, digested pMRT1119 and [sic] was hydrolysed by KpnI, purified with the aid of the "QIAquick PCR Purification" kit, recovered in 50 µl of H₂O, subjected to the action of 6 units of T4 DNA polymerase (New England Biolabs) in a reaction medium of 120 µl in the presence of 12 µl of T4 DNA polymerase x 10 buffer, 4 µl of 10 mM dNTPs and 6 µl of BSA at 1 mg/ml. The reaction was carried out at 37°C for 30 min. The vector fragment was then isolated by 0.8% agarose gel electrophoresis, purified with the aid of the "QIAquick Gel Extraction" kit and recovered in 50 µl of H₂O.

Thirty nanograms of the plasmid so treated were religated by ligation by PCR reaction (procedure described in 7).

The resultant plasmid was called pMRT1191. It is represented in Figure 10 and its complete sequence SEQ.ID10 is given in the sequence listing.

10.1.2. Obtaining pMRT1201

The plasmid pMRT1201 (7943 bp) differs from pMRT1191 by the insertion of the expression cassette of the uidA gene isolated from the plasmid pUC19-Phmwig-IA-uidA-Tnos which comprises the uidA expression cassette inserted into the EcoRI and HindIII sites of pUC19

(marketed by New England Biolabs). This plasmid results from successive clonings. The fragment corresponding to the sequence «uidA gene - nos terminator» was isolated by successive digestions by «EcoRI followed by the action of the Klenow fragment» and SmaI from pBI221 SacI (= pBI221 (Clontech) whose ScaI site was eliminated by SacI digestion followed by the action of T4 DNA polymerase, then ligation). The isolated and purified fragment was inserted into the «SmaI – [SphI + T4 DNA polymerase]» sites of pUC19 (Clontech) in order to obtain the resultant plasmid pUC19-uidA-Tnos. After that, the EcoRI site recreated on the 3' side of Tnos was eliminated by replacement of the BstBI-HindIII fragment of pUC19-uidA-Tnos with the BstBI-HindIII fragment obtained by digestion of the fragment amplified by PCR (1054 bp) from the pUC19-uidA-Tnos matrix with the aid of the 2 oligodeoxynucleotides 5' AGGCATTGGTTTCGAAGCG 3' (SEQ.ID46) containing the BstBI site and 5' TACGCCAAGCTTGGCAATTCC 3' (SEQ.ID47). The resultant plasmid was called pUC19-uidA-Tnos EcoRI. Then, in order to introduce a NcoI site at the level of the initiator codon of the uidA gene, a 440 bp fragment was amplified by PCR from the pUC19-uidA-Tnos EcoRI matrix with the aid of the 2 oligodeoxynucleotides 5' AATACCCGGGACCATGGTCCGTCCTGTAG 3' (SEQ.ID48) containing the NcoI and SmaI sites and 5' ATAGTCTGCCAGTTCAGTTCGTTG 3' (SEQ.ID49) situated downstream of SnaBI. The PCR fragment digested by SmaI and SnaBI was then inserted into pUC19-uidA-Tnos EcoRI in order to replace the existing SmaI-SnaBI fragment. The resultant plasmid was called pUC19-NcoI-uidA-Tnos. After that, the Phmwg promoter [High Molecular Weight Glutenin, Anderson et al. (1989)], carried by the EcoRI - SmaI fragment, was introduced into the EcoRI and SmaI sites of pUC19-NcoI-uidA-Tnos in order to produce the plasmid pUC19-Phmwg-uidA-Tnos. Finally, the intron 1 of the rice actin gene (IA, EcoRV – SmaI fragment) comes from the plasmid pAct1-F6 (McElroy D et al., 1991) which is itself modified by the addition of restriction sites notably on both sides of the EcoRV-SmaI fragment containing IA. The SmaI - NcoI fragment carrying IA was isolated from pAct1-F6 modified and inserted into the SmaI and NcoI sites of pUC19-Phmwg-uidA-Tnos in order to produce the plasmid pUC19-Phmwg-IA-uidA-Tnos. The techniques used are the ones already described in this patent.

The pMRT1191 vector fragment was obtained by digestion of 2 µg of pMRT1191 by EcoRI, purification with the aid of the «Concert Rapid PCR Purification System» kit, followed

by the action of 20 units of the Klenow fragment (New England Biolabs) in the presence of 12 µl of 500 mM Tris-HCl pH 7.5, 500 mM MgCl₂, 6 µl of 1M dithiothreitol, 6 µl of each of the dNTPs at 10 mM in a reaction volume of 120 µl at 37°C for 30 min., purification with the aid of the «Concert Rapid PCR Purification System» kit, and recovery in 50 µl of H₂O. After that,
5 digested and treated pMRT1191 was dephosphorylated with 50 units of calf intestinal alkaline phosphatase (New England Biolabs) in a final reaction medium of 120 µl in the presence of 12 µl of 3 x 10 buffer (New England Biolabs) at 37°C for 1 hour, purified with the aid of the «Concert Rapid PCR Purification System» kit, and recovered in 50 µl of H₂O.

The insert DNA fragments (2 µg) corresponding to the expression cassette of the uidA
10 gene (3 kbp), viz. the uidA gene placed under the control of the wheat "high molecular weight glutenin" promoter followed by the intron 1 of the rice actin gene and the nopaline synthase terminator of Agrobacterium tumefaciens, were obtained by digestion of pUC19-Phmwg-IA-uidA-Tnos by ScaI (site present in pUC19), purification with the aid of the "QIAquick PCR Purification" kit, recovery in 50 µl of H₂O, digestion by HindIII, purification with the aid of the
15 "QIAquick PCR Purification" kit, recovery in 50 µl of H₂O, treatment by 20 units of the Klenow fragment (New England Biolabs) in the presence of 12 µl of 500 mM Tris-HCl, pH 7.5, 500 mM MgCl₂, 6 µl of 1M dithiothreitol, 6 µl of each of the 10 mM dNTPs in a reaction volume of 120 µl at 37°C for 30 min., purification with the aid of the "QIAquick PCR Purification" kit, recovery in 50 µl of H₂O and digestion by EcoRI. After that, the insert DNA fragments were
20 isolated by 0.8% agarose gel electrophoresis, purified with the aid of the "QIAquick Gel Extraction" kit, subjected to the action of the Klenow fragment as described above, and recovered in 50 µl of H₂O.

The ligation by PCR reaction was carried out as described earlier in 7, with 10 ng of dephosphorylated digested pMRT1191 plasmid and 100 ng of digested and treated insert DNA
25 fragments. The bacteria, Escherichia coli DH5⁺, which had previously been made competent, were transformed. The plasmid DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method and verified by enzymatic digestions and sequencing. The resultant plasmid was called pMRT1201 (7943 bp).

It is represented in Figure 15 and its complete sequence SEQ.ID15 is given in the sequence listing.

10.1.3. Obtaining pMRT1210

The plasmid pMRT1210 (10003 bp) differs from pMRT1201 by the insertion of the sequence "promoter followed by the intron 1 of the rice actin gene - bar gene" (Pact-IA-bar) isolated from pSB12-Pact-IA-bar-Tnos which derives from pSB12 described by Komari et al. (1996). The plasmid pSB12-Pact-IA-bar-Tnos results from the cloning of the expression cassette «Pact-IA-bar-Tnos» (BspDI - «XhoI + Klenow» fragment), isolated from pDM302, into the SmaI and BspDI sites of pSB12, whose XhoI site on the LB side was deleted. The plasmid pDM302, constructed in the laboratory of Wu R., comprises the expression cassette «Pact-IA-bar-Tnos» in the plasmid pSP72 marketed by Promega.

The pMRT1201 vector fragment was obtained by digestion of 2 µg of pMRT1201 by HpaI, purification with the aid of the «Concert Rapid PCR Purification System» kit, and recovery in 98 µl of H₂O. After that, digested pMRT1201 was dephosphorylated with 50 units of calf intestinal alkaline phosphatase (New England Biolabs) in a final reaction medium of 120 µl in the presence of 12 µl of 3 x 10 buffer (New England Biolabs) at 37°C for 1 hour, purified with the aid of the «Concert Rapid PCR Purification System» kit, and recovered in 50 µl of H₂O.

The insert DNA fragments (2 µg) corresponding to "Pact-IA-bar" (2.1 kbp) were obtained by XbaI digestion of pBIOS273, purification with the aid of the "QIAquick PCR Purification" kit, recovery in 50 µl of H₂O, treatment with 20 units of the Klenow fragment (New England Biolabs) in the presence of 12 µl of 500 mM Tris-HCl, pH 7.5 - 500 mM MgCl₂, 6 µl of 1M dithiothreitol and 6 µl of each of the 10 mM dNTPs in a reaction volume of 120 µl at 37°C for 30 min. The insert DNA fragments so treated were then isolated by 0.8% agarose gel electrophoresis, purified with the aid of the "QIAquick PCR Purification" kit and recovered in 50 µl of H₂O.

The ligation by PCR reaction was carried out as described earlier in 7, with 60 ng of dephosphorylated digested pMRT1201 plasmid and 100 ng of digested and treated insert DNA fragments. The bacteria, *Escherichia coli* DH5⁺, which had previously been made competent,

were transformed. The plasmid DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method and verified by enzymatic digestions and sequencing. The resultant plasmid was called pMRT1210. It is represented in Figure 21 and its complete sequence SEQ.ID21 is given in the sequence listing.

5 This plasmid pMRT1210 was then introduced into *Agrobacterium tumefaciens* strain LBA4404 (pSB1) [Komari T et al., 1996] and strain LBA4404 by direct transformation in accordance with the procedure of Holsters et al. (1978).

10.1.4. Obtaining pMRT1193

10 The binary plasmid pMRT1193 was obtained by cloning, into the EcoRI and [XhoI + Klenow fragment action] sites of pMRT1119, of the «EcoRI – [HindIII + Klenow fragment action]» fragment carrying the sequence «Phmwg-IA-uidA-Tnos» isolated from pUC19-Phmwg-IA-uidA-Tnos, described in 10.1.2.

15 To do this, the pMRT1119 vector fragment was obtained by digestion of 10 µg of pMRT1119 by XhoI, purification with the aid of the "QIAquick PCR Purification" kit, recovery in 50 µl of H₂O, treatment with 20 units of the Klenow fragment (New England Biolabs) in the presence of 12 µl of 500 mM Tris-HCl, pH 7.5 - 500 mM MgCl₂, 6 µl of 1M dithiothreitol and 6 µl of each of the 10 mM dNTPs in a reaction volume of 120 µl at 37°C for 30 min. Then purification with the aid of the "QIAquick PCR Purification" kit and digestion by EcoRI. The treated vector fragment was isolated by 0.8% agarose gel electrophoresis, purified with the aid of the "QIAquick PCR Purification" kit and recovered in 50 µl of H₂O. After that, digested pMRT1119 was dephosphorylated with 50 units of calf intestinal alkaline phosphatase (New England Biolabs) in a final reaction medium of 120 µl in the presence of 12 µl of 3 x 10 buffer (New England Biolabs) at 37°C for 1 hour, purified with the aid of the "QIAquick PCR Purification" kit and recovered in 50 µl of H₂O.

25 The insert DNA fragments (2 µg) corresponding to the expression cassette of the uidA gene (3 kbp), viz. the uidA gene placed under the control of the wheat "high molecular weight glutenin" promoter followed by the intron 1 of the rice actin gene and the nopaline synthase terminator of *Agrobacterium tumefaciens*, were obtained by digestion of pUC19-Phmwg-IA-

uidA-Tnos by ScaI (site present in pUC19), purification with the aid of the "QIAquick PCR Purification" kit, recovery in 50 µl of H₂O, digestion by HindIII, purification with the aid of the "QIAquick PCR Purification" kit, recovery in 50 µl of H₂O, treatment with 20 units of the Klenow fragment (New England Biolabs) in the presence of 12 µl of 500 mM Tris-HCl, pH 7.5, 500 mM MgCl₂, 6 µl of 1M dithiothreitol and 6 µl of each of the 10 mM dNTPs in a reaction volume of 120 µl at 37°C for 30 min., purification with the aid of the "QIAquick PCR Purification" kit, recovery in 50 µl of H₂O, and digestion by EcoRI. The insert DNA fragments were then isolated by 0.8% agarose gel electrophoresis, purified with the aid of the "QIAquick Gel Extraction" kit and recovered in 50 µl of H₂O.

The ligation by PCR reaction was carried out as described earlier in 7, with 20 ng of dephosphorylated digested pMRT1119 plasmid and 100 ng of digested and treated insert DNA fragments. The bacteria, *Escherichia coli* DH5⁺, which had previously been made competent, were transformed. The plasmid DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method and verified by enzymatic digestions and sequencing. The resultant plasmid was called pMRT1193 (9143 bp). It is represented in Figure 12 and its complete sequence SEQ.ID12 is given in the sequence listing.

10.2. Obtaining pMRT1195

The plasmid pMRT1195 (6857 bp) differs from pMRT1191 by the insertion of the sequence "promoter followed by the intron 1 of the rice actin gene - bar gene" (Pact-IA-bar). The vector pMRT1191 was digested by Bsp120I and KpnI followed by the action of the enzyme T4 DNA polymerase and dephosphorylated but not religated was obtained as described in 10.1.1.

The insert DNA fragments corresponding to "Pact-IA-bar" (2.1 kbp) were obtained as described in 10.1.3.

The ligation by PCR reaction was carried out as described earlier in 7, with 20 ng of dephosphorylated digested pMRT1191 plasmid and 100 ng of digested and treated insert DNA fragments. The bacteria, *Escherichia coli* DH5a, which had previously been made competent, were transformed. The plasmid DNA of the clones obtained, selected on LB medium

supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method and verified by enzymatic digestions and sequencing. The resultant plasmid was called pMRT1195. It is represented in Figure 13 and its complete sequence SEQ.ID13 is given in the sequence listing.

10.3. Obtaining pMRT1202

5 The plasmid pMRT1202 (5614 bp) differs from pMRT1195 by the replacement of the sequence "Pact-IA" by the nopaline synthase (Pnos) promoter of *Agrobacterium tumefaciens* isolated from pMRT1121 produced in *Escherichia coli* strain SCS110.

10 The pMRT1195 vector fragment (2 µg) was digested by PstI, purified with the aid of the «Concert Rapid PCR Purification System» kit, recovered in 98 µl of H₂O, treated by the action of 6 units of T4 DNA polymerase (New England Biolabs) in a reaction medium of 120 µl in the presence of 12 µl of T4 DNA polymerase x 10 buffer, 4 µl of 10 mM dNTPs and 6 µl of BSA at 1 mg/ml. The reaction was carried out at 37°C for 30 min. The vector fragment was then purified with the aid of the «Concert Rapid PCR Purification System» kit, recovered in 50 µl of H₂O, digested by HindIII, isolated by 0.8% agarose gel electrophoresis, purified with the aid of the «Concert Rapid PCR Purification System» kit, and recovered in 98 µl of H₂O. After that, digested and treated pMRT1195 was dephosphorylated with 50 units of calf intestinal alkaline phosphatase (New England Biolabs) in a final reaction medium of 120 µl in the presence of 12 µl of 3 x 10 buffer (New England Biolabs) at 37°C for 1 hour, purified with the aid of the «Concert Rapid PCR Purification System» kit, and recovered in 50 µl of H₂O.

20 The insert DNA fragments corresponding to "Pnos" were obtained by BspEI digestion of pMRT1121, purification with the aid of the «Concert Rapid PCR Purification System» kit, recovery in 98 µl of H₂O, treatment with 20 units of the Klenow fragment (New England Biolabs) in the presence of 12 µl of 500 mM Tris-HCl, pH 7.5 - 500 mM MgCl₂, 6 µl of 1M dithiothreitol and 6 µl of each of the 10 mM dNTPs in a reaction volume of 120 µl at 37°C for 30 min., purification with the aid of the «Concert Rapid PCR Purification System» kit, recovery in 50 µl of H₂O, and digestion by HindIII. The insert DNA fragments (219 bp) carrying Pnos (209 bp) were then isolated by 2% agarose gel electrophoresis, purified with the aid of the «Concert Rapid PCR Purification System» kit, and recovered in 50 µl of H₂O.

The ligation by PCR reaction was carried out as described earlier in 7, with 48 ng of dephosphorylated digested pMRT1195 plasmid and 56 ng of digested and treated insert DNA fragments. The bacteria, *Escherichia coli* DH5⁺, which had previously been made competent, were transformed. The plasmid DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method and verified by enzymatic digestions and sequencing. The resultant plasmid was called pMRT1202. It is represented in Figure 16 and its complete sequence SEQ.ID16 is given in the sequence listing.

10.4. Obtaining pMRT1212

The plasmid pMRT1212 (8987 bp) differs from pMRT1206 by the replacement of the nptII gene by the bar gene.

To do this, the pMRT1206 vector fragment was obtained by digestion by Bsu36I of 2 µg of pMRT1206, purification with the aid of the «Concert Rapid PCR Purification System» kit, recovery in 50 µl of H₂O and digestion by AatII. The digestion by Bsu36I and AatII allowed deletion of the vector of the fragment corresponding to «part Pnos nptII Tnos LB». The digested vector fragment was then subjected to 1% agarose gel electrophoresis, purified with the aid of the «Concert Rapid PCR Purification System» kit and recovered in 98 µl of H₂O. This digested and treated vector fragment was then dephosphorylated with 50 units of calf intestinal alkaline phosphatase (New England Biolabs) in a final reaction medium of 120 µl in the presence of 12 µl of 3 x 10 buffer (New England Biolabs) at 37°C for 1 hour, purified with the aid of the «Concert Rapid PCR Purification System» kit and recovered in 50 µl of H₂O.

The insert DNA fragments corresponding to «part Pnos bar Tnos LB» (1.2 kbp) were obtained by digestion by Bsu36I of 2 µg of pMRT1202, purification with the aid of the Concert Rapid PCR Purification System kit, recovery in 40 µl of H₂O, digestion by AatII, isolation by 1% agarose gel electrophoresis, purification with the aid of the Concert Rapid PCR Purification System kit and recovery in 50 µl of H₂O.

The ligation by PCR reaction was carried out as described earlier in 7, with 100 ng of dephosphorylated digested pMRT1206 plasmid and 50 ng of digested insert DNA fragments. The bacteria, *Escherichia coli* DH5⁺, which had previously been made competent, were transformed.

The plasmid DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method and verified by enzymatic digestions and sequencing. The resultant plasmid was called pMRT1212. It is represented in Figure 22 and its complete sequence SEQ.ID22 is given in the sequence listing.

11. Evaluation of the synthetic vectors produced in the bacteria

The yield of plasmids obtained from the bacteria *Escherichia coli* strain DH5 was determined.

I Non-binary synthetic plasmids

Primary cultures of the recombinant bacteria possessing the plasmids pMRT1105, pMRT1106 or pMRT1105-ori ColE1 were prepared from 500 ml of each of the respective "glycerol-stocks" in 10 ml of LB medium supplemented with kanamycin at 50 mg/l, at 37°C for 16 hours. After that, 500 ml of each of the primary cultures was cultured in 100 ml of LB supplemented with kanamycin at 50 mg/l at 37°C for 23 hours. For pMRT1106, two cultures were prepared, one without addition of chloramphenicol and the other with addition of chloramphenicol at 40 mg/l after 7 hours of culture. The cultures were then centrifuged at 5000 g for 10 min. The plasmid DNAs were extracted with the aid of the "QIAFilter Plasmid Midi kit" (QIAGEN), in accordance with the manufacturer's recommendations, and quantitatively determined with the spectrophotometer. The quantities of plasmids obtained were determined as 10 mg, 50.6 mg, 86.2 mg and 9.8 mg respectively for pMRT1105, pMRT1106, pMRT1106 treated with chloramphenicol, and pMRT1105 ori ColE1 (plasmid pMRT1105 in which ori ColE1 was inserted in the reverse orientation to that of pMRT1106). These results show that the synthetic plasmids are replicated in *Escherichia coli*, the presence of the origin of replication "ori ColE1" in the right orientation (NdeI site of "ori ColE1" near "ori RK2") allows the yield to be increased by a factor of 5, whereas no effect is shown for the reverse orientation of "ori ColE1", treatment with chloramphenicol contributes an additional factor of 1.6.

11.2. Synthetic binary plasmids.

The synthetic binary plasmids pMRT1118 and pMRT1119 are evaluated relative to the controls pBin19 and pBIOC4. The plasmid pBIOC4 differs from pGA492 (An,1986) by the deletion of virtually all the coding sequence of the pGA492 cat gene and by the conversion of the HindIII site into an EcoRI site. It was obtained by double digestion by SacI and ScaI followed by the action of the enzyme T4 DNA polymerase (New England Biolabs) and ligation of the modified plasmid (20 ng) in a reaction medium of 10 µl in the presence of 1 µl of T4 DNA ligase x 10 buffer (Amersham) and 2.5 units of T4 DNA ligase (Amersham) at 14°C for 16 hours. The bacteria, *Escherichia coli* strain DH5⁺, which had previously been made competent, were transformed (Hanahan, 1983). The plasmid DNA of the clones obtained, selected on tetracycline at 12 mg/l, was extracted by the alkaline lysis method (Birnboim and Doly, 1979) and analysed by enzymatic digestions. The HindIII site of the plasmid DNA of the clone selected were then modified into an EcoRI site with the aid of a phosphorylated HindIII-EcoRI adaptor (Stratagene). To do this, 500 ng of plasmid DNA of the clone selected were digested by HindIII, dephosphorylated by the enzyme calf intestinal alkaline phosphatase (Boehringer Mannheim) in accordance with the manufacturer's recommendations and coprecipitated in the presence of 1500 ng of HindIII-EcoRI adaptor DNA, 0.1 volume of 3M sodium acetate, pH 4.8, and 2.5 volumes of absolute ethanol at -80°C for 30 min. After centrifugation at 12000 g for 30 min., the precipitated DNA was washed with 70% ethanol, dried, recovered in 8 µl of H₂O, held at 65°C for 10 min., then ligated as described earlier. After inactivation of the T4 DNA ligase at 65°C for 10 min., the ligation reaction mixture was digested by EcoRI, purified by 0.8% agarose gel electrophoresis, electroeluted, precipitated with absolute ethanol, centrifuged at 12000 g for 30 min., washed with 70% ethanol, dried, then ligated and introduced into *Escherichia coli* strain DH5⁺, and treated as described earlier.

Primary cultures of the recombinant bacteria possessing the plasmids mentioned above were prepared from 500 ml of each of the respective "glycerol-stocks" in 10 ml of LB medium supplemented with kanamycin at 50 mg/l, except for pBIOC4, which was selected on tetracycline at 12 mg/l, at 37°C for 16 hours. After that, 250 ml of each of the primary cultures were cultured in 100 ml of LB supplemented with the appropriate antibiotic at 37°C for 23 hours. The cultures

were then centrifuged at 5000 g for 10 min. The plasmid DNAs were extracted with the aid of the "QIAfilter Plasmid Midi kit" (QIAGEN) in accordance with the manufacturer's recommendations and quantitatively determined with the spectrophotometer. The quantities of plasmids obtained were determined as 94.2 mg, 113 mg, 37.2 mg and 45.6 mg respectively for pMRT1118, pMRT1119, pBin19 and pBIOC4.

These results show that the synthetic binary plasmids are replicated in *Escherichia coli*, and that the yield of pMRT1119 is increased by the factors 3 and 2.5 respectively, relative to the pBin19 and pBIOC4 controls.

Furthermore, these synthetic plasmids are replicated in *Agrobacterium tumefaciens* strain LBA4404.

12. Evaluation of the synthetic plasmids by transgenesis

12.1. Transformation of tobacco

12.1.1. Stable transformation of tobacco

The transformation of tobacco (*Nicotiana tabacum* L. var. PBD6) was carried out by infecting foliar discs isolated from 6-week-old tobacco seedlings in vitro with recombinant agrobacteria in accordance with the method described by Horsch et al. (1985).

All the in vitro cultures are prepared in a controlled environment chamber in which the light intensity is 200 $\mu\text{E}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$, the photoperiod 16 hours light/8 hours darkness, and the temperature 25°C.

Except for the initial coculture stage, the regeneration, development and rooting stages were performed on different selective media supplemented with a bacteriostat, namely augmentin at 400 mg/l, and a selective agent, namely kanamycin at 200 or 100 mg/l.

The different stages and the media used are as follows:

After primary culture of the agrobacteria in 5 ml of 2YT medium (bacto-tryptone 10 g/l, yeast extract 5 g/l, NaCl 5 g/l, pH 7.2) supplemented with CaCl_2 , with a final concentration of

6 mM, and with appropriate antibiotics, at 28°C for 48 hours, a culture is performed in 10 ml of 2YT medium, supplemented with CaCl₂ and antibiotics, at 28°C for a period of one night. The culture is then centrifuged at 3000 g for 10 min. and the bacteria are resuspended in 10 ml of liquid MS30 (M0404 at 4.4 g/l, marketed by SIGMA, supplemented with sucrose, 30 g/l, pH 5.7).

The coculture is performed by placing the foliar explants cut from the leaves of the seedlings in vitro, about 1 cm², in contact with the suspension of agrobacteria, diluted 1/10, in liquid MS30 for 20 min. The explants so treated are then rapidly dried on filter paper and placed on a solid coculture medium (CM) (MS30, Benzyl Amino Purine (BAP) 1 mg/l, Indole-3 Acetic Acid (ANA) at 0.1 mg/l, agar at 8 g/l) for 48 hours in the controlled environment chamber.

The treated explants are then placed on a solid regeneration medium (solid CM, augmentin 400 at mg/l, kanamycin at 200 mg/l). The explants are pricked out on the same medium after 2 weeks.

After 2 weeks the shoots are pricked out on a solid development medium (M0404 at 4.4 g/l, marketed by SIGMA, supplemented with sucrose 20 g/l, pH 5.7 (liquid MS20), augmentin at 400 mg/l, kanamycin at 100 mg/l, agar at 8 g/l).

After 2 weeks, the transformed seedlings are pricked out on a solid rooting medium identical to the development medium. Rooting takes 2 to 3 weeks, after which the seedlings are removed to the growth chamber in Giffy pots for 10 days (photoperiod 16 hours light/8 hours darkness, 23°C, 70% relative humidity), then placed in the greenhouse.

12.1.2. Evaluation of the synthetic plasmids used in transformation of tobacco

The recombinant agrobacteria (*Agrobacterium tumefaciens* strain LBA4404) containing the synthetic binary plasmid pMRT1118 or the control binary plasmid pBin19 were used to transform the tobacco *Nicotiana tabacum* L. var. PBD6.

The results showed that the number and growth of the seedlings which develop in the presence of the selective agent (kanamycin) are similar for the two binary plasmids tested. These

observations indicate that the synthetic binary plasmid is perfectly functional in transgenesis of tobacco.

12.2. Transformation of maize

12.2.1. Genetic transformation of maize calli

5 The genetic transformation of maize, whatever method is employed (electroporation, Agrobacterium, microfibres, particle gun), generally requires the use of cells that are undifferentiated in rapid divisions, having retained the ability to regenerate whole plants. Cells of this type make up the friable embryogenic callus (called Type II) of maize.

10 These calli were obtained from immature embryos of genotype HI, II or (A188 x B73) in accordance with the method and on the media described by Armstrong (Maize handbook, 1994, Freeling M and Walbot V (eds), pp 665-671). The calli so obtained were multiplied and maintained by successive subcultures every 2 weeks on the starting medium.

15 Seedlings were regenerated from these calli by modifying the hormonal and osmotic equilibrium of the cells in accordance with the method described by Vain et al. (1989). These plants were then hardened off in the greenhouse, where they can be crossed or selfed.

12.2.2. Genetic transformation of maize with the particle gun

20 The production and regeneration of the cell lines needed for transformation are described in 12.2.1.

The procedure which uses the particle gun for genetic transformation is described by 20 Finer et al. (1992). The target cells are callus fragments with a surface area of 10 to 20 mm². These fragments were placed on a starting medium supplemented with 0.2 M mannitol and 0.2 M sorbitol for 4 hours before bombardment.

The tungsten particles (M10) and transformation plasmids were coprecipitated in accordance with the protocol described by Klein (1987). The particles thus coated were projected

towards the target cells by means of the particle gun in accordance with the protocol of Finer et al. (1992).

The bombarded calli were placed in darkness at 27°C. After 24 hours, the first subculturing took place, followed by subculturing every 2 weeks for 3 months on a starting medium to which the appropriate selective agent to select only the transformed calli had been added. These calli were then grown in the presence of the selective agent in order to regenerate transformed seedlings, as described in 12.2.1. The seedlings obtained were hardened off and transferred to the greenhouse, where they can be crossed or selfed.

12.2.3. Genetic transformation of maize by *Agrobacterium tumefaciens*

The technique used is described by Ishida et al. (1996). Immature embryos 1.0 to 1.2 mm in length (9 to 14 days after pollination) were washed in the LS-inf medium, then immersed in the suspension of agrobacteria, prepared as described by Ishida et al. (1996), vortexed for 30 sec., and incubated at ambient temperature for 5 min. The immature embryos so treated were cultured on LS-AS medium in darkness at 25°C for 3 days, then transferred to LSD 1.5 medium supplemented with phosphinothricin at 5 mg/l and cefotaxime at 250 mg/l, in darkness at 25°C for 2 weeks, and finally placed on LSD 1.5 medium supplemented with phosphinothricin at 10 mg/l and cefotaxime at 250 mg/l, in darkness at 25°C for 3 weeks. The Type I calli thus generated were isolated, fragmented and transferred to LSD 1.5 medium supplemented with phosphinothricin at 10 mg/l and cefotaxime at 250 mg/l, in darkness at 25°C for 3 weeks. The Type I calli which proliferated were then isolated and placed on LSZ medium supplemented with phosphinothricin at 5 mg/l and cefotaxime at 250 mg/l, subjected to a photoperiod of 16 hours light/8 hours darkness at 25°C for 2 to 3 weeks. The regenerated seedlings were then transferred to ½ LSF medium subjected to a photoperiod of 16 hours light/8 hours darkness at 25°C for 1 to 2 weeks, and then transferred to the growth chamber and greenhouse.

13. Influence of the vector backbone on gene expression

Stably transformed tobacco plants were obtained for the synthetic binary plasmids pMRT1204 bearing the wild-type nptII gene (16 plants) and pMRT1206 bearing the mutated nptII gene (12 plants), and also for the control vector pBI121 sold by Clontech which

corresponds to plasmid pBIN19 bearing the wild-type nptII gene and containing the expression cassette “P35S – uidA – polyA nos” (11 plants).

The influence of the synthetic vector backbone and that of the control was evaluated by ELISA analysis of protein extracts using the NPTII ELISA kit sold by 5 Prime -> 3 Prime. The analysis enabled a quantification of the NPTII protein produced (in micrograms, µg) with respect to the amount of soluble extracted protein (in milligrams, mg).

The proteins were extracted in “Tris-HCl 25 mM pH7.8 ; Phenylmethylsulfonylfluoride 1 mM” buffer from tobacco leaves ground in liquid nitrogen. After centrifugation for 10 minutes at 10000 g and 4°C, the extracted soluble proteins contained in the supernatant were measured according to Bradford's method (A rapid and sensitive method for the detection of microgram quantities of proteins utilizing the principle of protein-dye binding. Anal. Biochem. (1976)72, 248-254).

The antibodies used for ELISA quantification were rabbit polyclonal antibodies specifically recognizing the coating NPTII protein and biotinylated anti-NPTII antibodies for the detection of the NPTII protein.

The results of the analysis shown in Figure 23 indicate that :

⑩ for the wild type nptII gene cassette and different plasmid backbones, the quantities of NPTII obtained with the synthetic binary plasmid pMRT1204 are greater by a factor of 6.7 than those obtained with the original pBIN19 binary plasmid, which demonstrates the efficiency of the synthetic vector.

⑩ for an identical plasmid backbone, i.e. synthetic binary plasmids pMRT1204 and pMRT1206, but with different nptII genes, i.e. Wild type and mutated nptII, the quantities of NPTII obtained in the wild type gene differ from those obtained for the mutated nptII gene by a factor 1.2.

14. Synthesis of pMRT1334.

The plasmid pMRT1334 (9688 bp) was obtained by replacing the nptII expression

cassette of pMRT1206 by the nptII expression cassette of pBIN19.

The vector fragment derived from pMRT1206 was obtained by digestion of 10 µg of pMRT1206 with KpnI, purified using a “Concert Rapid PCR Purification System” kit, and taken up in 50 µl of water. Then the digested pMRT1206 was subjected to the action of 6 units of T4 DNA polymerase (New England Biolabs) in a reaction mixture of 120 µl in the presence of 12 µl of T4 10x DNA polymérase buffer, 4 µl of 10 mM dNTP and 6 µl of 1 mg/ml BSA. The reaction was carried out at 37°C for 30 minutes. The digested and thus treated pMRT1206 vector was purified with a “Concert Rapid PCR Purification System” kit, taken up in 50 µl of water, then digested with AflII. The vector fragment derived from pMRT1206 was isolated by electrophoresis on a 1% agarose gel, purified with a “Concert Rapid PCR Purification System” and taken up in 50 µl of water.

The DNA insert fragment DraI – AflII (1.5 kbp), corresponding to the nptII expression cassette was obtained by digestion of 9 µg of pBIN19 plasmid with DraI and AflII. Then, the fragment was subjected to electrophoresis on a 1% agarose gel in TEB buffer, purified using a “Concert Rapid PCR Purification System”, and taken up in 50 µl of water.

The PCR ligation reaction was carried out using 100 ng of the plasmid fragment derived from pMRT1206 and 100 ng of the DNA insert fragment DraI – AflII in a reaction mixture of 20 µl in the presence of 2 µl of T4 10x DNA ligase buffer (Epicentre Technologies), 2 µl 2.5 mM ATP and 4 units of T4 DNA ligase (Epicentre Technologies). The reaction consists of 180 cycles each including 2 steps, the first at 10°C for 30 seconds and the second at 30°C for 30 seconds in a "GeneAmp PCR System 9700" thermocycler.

Previously prepared competent *Escherichia coli* DH5 bacteria, were transformed (Hanahan, 1983). The plasmid DNA of the obtained clones, selected on LB media supplemented with kanamycin (50 mg/l), was extracted according to the alkaline lysis method and verified by enzymatic digestion and sequencing. The resulting plasmid was designated pMRT1334 (Figure 24). Its complete sequence is given under SEQ.ID50 in the sequence listing.

15. Synthesis of pMRT1335

The plasmid pMRT1335 (15208 bp) is a control vector and results from the insertion of the expression cassette “ep35S-gus (uidA)-polyA35S” isolated from pMRT1206 into pBIN19.

13 µg of pBIN19 plasmid were digested with EcoRI, purified using a “Concert Rapid PCR Purification System”, and taken up in 50 µl water. Then the previously digested pBIN19 was subjected to the action of 20 units of Klenow fragment (New England Biolabs) in the presence of 12 µl 500 mM pH7.5 Tris-HCl 500 mM MgCl₂, 6 µl of 1M dithiothreitol, 6 µl each of 10 mM dNTP in a reaction volume of 120 µl at 37°C for 60 minutes. The digested pBIN19 vector was purified using a “Concert Rapid PCR Purification System”, taken up in 50 µl water, then digested with KpnI. The vector fragment thus produced was isolated by electrophoresis on a 1% agarose gel, and purified using a “Concert Rapid PCR Purification System” and taken up in 50 µl of water.

The DNA insert fragment corresponding to the “ep35S-gus-polyA35S” (3.5 kbp) expression cassette was obtained by digestion of 10 µg of plasmid pMRT1206 with XhoI, purified using a “Concert Rapid PCR Purification System”, and taken up in 50 µl water, followed by the action of 20 units of Klenow fragment (New England Biolabs) in the presence of 12 µl of 500 mM pH7.5 Tris-HCl 500 mM MgCl₂, 6 µl 1M dithiothreitol, 6 µl each of 10 mM dNTP in a reaction volume of 120 µl at 37°C for 60 minutes, then purified using a “Concert Rapid PCR Purification System”, taken up in 50 µl water, and finally digested with KpnI. The DNA insert fragment thus produced was isolated by electrophoresis on a 1% agarose gel in TEB buffer, and purified using a “Concert Rapid PCR Purification System”, and taken up in 50 µl water.

The PCR ligation reaction was carried out with 100 ng of pBIN19 treated and 100 ng of DNA insert fragment as prepared previously in a reaction volume of 20 µl in the presence of 2 µl of T4 10x DNA ligase buffer (Epicentre Technologies), 2 µl 2.5 mM ATP and 4 units T4 DNA ligase (Epicentre Technologies). The reaction consists of 180 cycles each including 2 steps, the first at 10°C for 30 seconds and the second at 30°C for 30 seconds in a "GeneAmp PCR System 9700" thermocycler.

Previously prepared competent *Escherichia coli* DH5 bacteria, were transformed

(Hanahan, 1983). The plasmid DNA of the obtained clones, selected on LB media supplemented with kanamycin (50 mg/l), was extracted according to the alkaline lysis method and verified by enzymatic digestion and sequencing. The resulting plasmid was designated pMRT1335 (Figure 25). Its complete sequence is given under SEQ.ID51 in the sequence listing.

16. Synthesis of pMRT1336

The plasmid pMRT1336 (9285 bp) results from the insertion into pMRT1196 of the promoter MPr1165 (610 bp) isolated from plasmid pMRT1322 as described in PCT patent application PCT/IB00/00370, and incorporated into the present description by reference thereto for that relevant part.

10 $10\text{ }\mu\text{g}$ of pMRT1196 plasmid were digested with KpnI, purified using a "Concert Rapid PCR Purification System", taken up in $50\text{ }\mu\text{l}$ water, and redigested with HpaI. The thus treated vector fragment was subjected to electrophoresis on a 1% agarose gel, and purified using a "Concert Rapid PCR Purification System" and then taken up in $50\text{ }\mu\text{l}$ water.

15 The DNA insert fragment bearing the promoter MPr1165 (0.5 kbp) was obtained from $10\text{ }\mu\text{g}$ of pMRT1322 plasmid digested with KpnI, purified using a "Concert Rapid PCR Purification System" and taken up in $50\text{ }\mu\text{l}$ water, and then redigested with HpaI. The DNA insert fragment thus produced was subjected to gel electrophoresis on a 1% agarose gel in TEB buffer, purified using a "Concert Rapid PCR Purification System", and taken up in $50\text{ }\mu\text{l}$ water.

20 The PCR ligation reaction was carried out using 100 ng of vector fragment and 100 ng DNA insert fragment bearing the promoter MPr1165 as prepared above in a reaction volume of $20\text{ }\mu\text{l}$ in the presence of $2\text{ }\mu\text{l}$ T4 10x DNA ligase buffer (Epicentre Technologies), $2\text{ }\mu\text{l}$ 2.5 mM ATP and 4 units T4 DNA ligase (Epicentre Technologies). The reaction consists of 180 cycles each including 2 steps, the first at 10°C for 30 seconds and the second at 30°C for 30 seconds in a "GeneAmp PCR System 9700" thermocycler.

25 Previously prepared competent *Escherichia coli* DH5 bacteria, were transformed (Hanahan, 1983). The plasmid DNA of the obtained clones, selected on LB media supplemented with kanamycin (50 mg/l), was extracted according to the alkaline lysis method and verified by

enzymatic digestion and sequencing. The resulting plasmid was designated pMRT1336 (Figure 26). Its complete sequence is given in the sequence listing under SEQ.ID52.

Plasmid pMRT1322 was obtained by the insertion of promoter MPr1165 into pMRT1176. 10 µg of pMRT1176 plasmid was digested with XbaI and EcoRI, purified using a “Concert Rapid PCR Purification System” kit, taken up in 50 µl water, dephosphorylated with 50 units of calf intestine alkaline phosphatase (New England Biolabs) in a final reaction volume of 120 µl in the presence 12 µl of 3x10 buffer (New England Biolabs) at 37 °C for 1 hour, and purified using a “Concert Rapid PCR Purification System” kit, and then taken up in 50 µl water. The DNA insert fragment corresponding to promoter MPr1165 was obtained by PCR amplification from 10 ng of pMRT1240 matrix DNA using 20 pmoles each of two oligodesoxynucleotides, 5' AGCTCTAGAGCTGCCTGCAGCACTAGTATCC 3'(SEQ.ID53) bearing the site XbaI and 5' CGGAATTCGGCCTCTAGGTTGTTGTGTTG 3'(SEQ.ID54) bearing the site EcoRI, using the enzyme Platinum Taq polymerase High Fidelity (GIBCO BRL Life Technologies) and following the suppliers instructions. The PCR amplification reaction was carried out in a "GeneAmp PCR System 9700" thermocycler. After denaturation at 94°C for 2 minutes, the DNA was subjected to 25 cycles, each including denaturation steps at 94°C for 45 seconds, hybridisation at 55°C for 45 seconds and elongation at 68°C for 45 seconds. On the final cycle the elongation was continued at 68°C for 3 minutes. The PCR product obtained was isolated by electrophoresis on a 1.5% agarose gel in TBE buffer and purified using a “Concert Rapid PCR Purification System” kit, and then taken up in 50 µl water. The PCR product was then digested with XbaI and EcoRI, purified using a “Concert Rapid PCR Purification System” kit, and taken up in 50 µl water. The PCR ligation reaction was carried out as described previously with 80 ng of pMRT1176 vector fragment and 110 ng of DNA insert fragment bearing the MPr1165 promoter. Previously prepared competent Escherichia coli DH5 bacteria, were transformed (Hanahan, 1983). The plasmid DNA of the obtained clones, selected on LB media supplemented with kanamycin (50 mg/l), was extracted according to the alkaline lysis method and verified by enzymatic digestion and sequencing. The resulting plasmid was designated pMRT1322.

Plasmid pMRT1240 was obtained by insertion of a DNA fragment into plasmid

pMRT1234, by treatment with Klenow fragment (New England Biolabs) according to the supplier's instructions of the HindIII sites and BamHI. The DNA insert fragment was the L5 promoter fragment obtained by double PstI digestion, and BamHI digestion, followed by the action of T4 DNA polymerase (New England Biolabs) according to the supplier's instructions, of plasmid pMRT1165, as described in PCT patent application PCT/IB00/00370, and incorporated into the present description by reference thereto for that relevant part.

Plasmid pMRT1234 was obtained by the insertion at the EcoRI site of pMRT1175 of a DNA insert fragment EcoRI corresponding to the cDNA coding for human lactoferrin isolated from the plasmid pHMWG-IA-PSSp-Lf, as described in previously published patent application WO98/50543.

17. Synthesis of pMRT1337

Plasmid pMRT1337 (8289 bp) was obtained from the insertion, into pMRT1205, of the gfp gene isolated from pBINm-gfp5-ER and described in Haseloff J et Siemering KR. 1998, "The uses of green fluorescent protein in plants", in Green fluorescent protein : Strategies, applications and protocols (Chalfie M et Kain S, eds, Wiley, pp 191-220).

10 µg of plasmid pMRT1205 was digested with EcoRI, purified using a "Concert Rapid PCR Purification System » kit, taken up in 50 µl water, subjected to the action of 20 units of Klenow fragment (New England Biolabs) in the presence of 12 µl of 500 mM pH7.5 Tris-HCl 500 mM MgCl₂, 6 µl 1M dithiothreitol, 6 µl each of 10 mM dNTP in a reaction volume of 120 µl at 37°C for 60 minutes, purified using a "Concert Rapid PCR Purification System" kit, taken up in 50 µl water, and then digested with BamHI. The thus treated vector fragment was subjected to electrophoresis on 1% agarose gel, purified using a "Concert Rapid PCR Purification System" kit and taken up in 50 µl water.

The DNA insert fragment bearing the gfp gene (0.7 kbp) was obtained from 10 µg of plasmid pBINm-gfp5-ER digested with SacI, purified using a "Concert Rapid PCR Purification System" kit, taken up in 50 µl water, and subjected to the action of 6 units of T4 DNA polymerase (New England Biolabs) in a reaction volume of 120 µl in the presence of 12 µl T4 10x DNA polymerase buffer, 4 µl 10 mM dNTP and 6 µl 1 mg/ml BSA. The reaction was

carried out at 37°C for 30 minutes. The thus treated plasmid pBINm-gfp5-ER was then purified using a “Concert Rapid PCR Purification System” kit, taken up in 50 µl water, and digested with BamHI. The DNA insert fragment thus produced was subjected to gel electrophoresis on 1% gel agarose in TEB buffer, purified using a “Concert Rapid PCR Purification System” kit and taken
5 up in 50 µl in water.

The PCR ligation reaction was carried out with 100 ng of pMRT1205 vector fragment and 100 ng DNA insert fragment bearing the gfp gene as prepared above in a reaction volume of 20 µl in the presence of 2 µl T4 10x DNA ligase buffer (Epicentre Technologies), 2 µl 2.5 mM ATP and 4 units T4 DNA ligase (Epicentre Technologies). The reaction comprised 180 cycles
10 each including two steps, the first at 10°C for 30 seconds, the second at 30°C for 30 seconds, in a "GeneAmp PCR System 9700" thermocycler.

Previously prepared competent Escherichia coli DH5⁺ bacteria, were transformed (Hanahan, 1983). The plasmid DNA of the obtained clones, selected on LB media supplemented with kanamycin (50 mg/l), was extracted according to the alkaline lysis method and verified by enzymatic digestion and sequencing. The resulting plasmid was designated pMRT1337 (Figure 27). Its complete sequence is given in the sequence listing under SEQ.ID55.

18. Synthesis of pMRT1341

The plasmid pMRT1341 (14108 bp) results from the replacement of the “ep35S-gus-polyA35S” expression cassette from pMRT1335 by the expression cassette “ep35S-gfp-polyA35S” isolated from pMRT1337.
20

10 µg of plasmid pMRT1335 was digested by AgeI and KpnI. The thus digested vector fragment was isolated by electrophoresis on a 0.8% agarose gel, purified using a “Concert Rapid PCR Purification System” kit and then take up in 50 µl water.

The DNA insert fragment corresponding to the expression cassette “ep35S-gfp-polyA35S” (2.4 kbp) was obtained by digesting with XhoI of 10 µg plasmid pMRT1337, purifying using a “Concert Rapid PCR Purification System” kit, taking up in 50 µl water, and the action of 20 units of Klenow fragment (New England Biolabs) in the presence of 12 µl 500
25

mM pH7.5 Tris-HCl 500 mM MgCl₂, 6 µl 1M dithiothreitol, 6 µl each of 10 mM dNTP in a reaction volume of 120 µl at 37°C for 30 minutes, then, purifying using a “Concert Rapid PCR Purification System” kit, taking up in 50 µl water, and finally digesting with KpnI. The thus preapred DNA insert fragment was isolated by electrophoresis on 0.8% agarose gel in TEB buffer, purified using a “Concert Rapid PCR Purification System” kit, taken up in 50 µl water, digested with AgeI, purified using a “Concert Rapid PCR Purification System” and taken up again in 50 µl water.

The PCR ligation reaction was carried out on 100 ng vector fragment derived from pMRT1335 et 100 ng DNA insert fragment as produced above in a reaction volume of 20 µl in the presence of 2 µl T4 10x DNA ligase buffer (Epicentre Technologies), 2 µl 2,5 mM ATP and 4 units of T4 DNA ligase (Epicentre Technologies). The reaction comprised 180 cycles each including 2 steps, the first at 10°C for 30 seconds, the second at 30°C for 30 seconds, in a "GeneAmp PCR System 9700" thermocycler.

Previously prepared competent *Escherichia coli* DH5 bacteria, were transformed (Hanahan, 1983). The plasmid DNA of the obtained clones, selected on LB media supplemented with kanamycin (50 mg/l), was extracted according to the alkaline lysis method and verified by enzymatic digestion and sequencing. The resulting plasmid was designated pMRT1341 (Figure 28). Its complete sequence is given in the sequence listing under SEQ.ID56.

19. Synthèse de pMRT1342

Plasmid pMRT1342 (15077 bp) results from the replacement of the expression cassette “ep35S-gus-polyA35S” from pMRT1335 by the expression cassette “L5-gus-polyA35S” isolated from pMRT1336.

10 µg of plasmid pMRT1335 was digested with AgeI and KpnI. The thus digested vector fragment was isolated by electrophoresis on 0.8% agarose gel, purified using a “Concert Rapid PCR Purification System” kit and taken up in 50 µl water.

The DNA insert fragment corresponding to the expression cassette “L5-gus-polyA35S” (3.4 kbp) was obtained by digestion with XhoI of 10 µg plasmid pMRT1336, purified using a

“Concert Rapid PCR Purification System” kit, taken up in 50 µl water, subjecting this to the action of 20 units Klenow fragment (New England Biolabs) in the presence of 12 µl 500 mM pH7.5 Tris-HCl 500 mM MgCl₂, 6 µl 1M dithiothreitol, 6 µl each of 10 mM dNTP in a reaction volume of 120 µl at 37°C for 30 minutes, then, purified using a “Concert Rapid PCR Purification System” kit, and taken up in 50 µl water, and finally digested with KpnI. The thus prepared DNA insert fragment was isolated by electrophoresis on a 0.8% agarose gel in TEB buffer, purified using a “Concert Rapid PCR Purification System” kit, taken up in 50 µl water, digested with AgeI, purified using a kit as above and taken up again in 50 µl water.

The PCR ligation reaction was carried out with 100 ng of vector fragment derived from pMRT1335 and 100 ng of DNA insert fragment prepared as described above in reaction mixture of 20 µl in the presence of 2 µl T4 10x DNA ligase buffer (Epicentre Technologies), 2 µl 2.5 mM ATP and 4 units T4 DNA ligase (Epicentre Technologies). The reaction is comprised of 180 cycles each including 2 steps, the first at 10°C for 30 seconds and the second at 30°C for 30 seconds in a "GeneAmp PCR System 9700" thermocycler.

Previously prepared competent *Escherichia coli* DH5⁺ bacteria, were transformed (Hanahan, 1983). The plasmid DNA of the obtained clones, selected on LB media supplemented with kanamycin (50 mg/l), was extracted according to the alkaline lysis method and verified by enzymatic digestion and sequencing. The resulting plasmid was designated pMRT1342 (Figure 29). Its complete sequence is given in the sequence listing under SEQ.ID57.

20. Production of recombinant agrobacteria

The binary plasmids were transferred into the *Agrobacterium tumefaciens* LBA4404 strain according to the technique described by Holsters et al. [Holsters M., Dewaele D., Depicker A., Messens E., Van Montagu M. et Schell J. (1978). Transfection and transformation of *Agrobacterium tumefaciens*. Mol. Gen. Genet. 136, 181-187.]. The plasmid DNA of the obtained clones, selected on LB media supplemented with rifampicin (50 mg/l), was extracted according to the alkaline lysis method, modified by the addition of lysozyme (25 mg/ml) in the cell resuspension buffer. The plasmid DNA obtained was analysed by enzymatic digestion. The agrobacteria clones obtained were used for plant genetic transformation.

References

An G. 1986. Development of plant promoter expression vector and their use for analysis of differential activity of nopaline synthase promoter in transformed tobacco cells. *Plant Physiol.* 81 :86-91.

5 Anderson OD, Green FC, Yip RE, Halford NG, Shewry PR and Malpica-Romero JM. 1989. Nucleotide sequences of the two high-molecular-weight glutenin genes from the D-genome of a hexaploid bread wheat, *Triticum aestivum* L. *Cheyenne. Nucl. Acids Res.* 17 :461-462.

Birnboim HC and Doly J. 1979. A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucl. Acids Res.* 7 :1513.

10 Depicker A, Stachel S, Dhaese P, Zambryski P and Goodman HM. 1982. Nopaline synthase : transcript mapping and DNA sequence. *J. Mol. Appl. Genet.* 1 :561-573.

Finer JJ, Vain P, Jones MW and McMullen MD. 1992. Development of the particle inflow gun for DNA delivery to plant cells. *Plant Cell Rep.* 11 :323-328.

15 Franck A, Guilley H, Jonard G, Richards K and Hirth L. 1980. Nucleotide sequence of cauliflower mosaic virus DNA. *Cell* 21 :285-294.

20 Frisch DA, Harris-Haller LW, Yokubaitis NT, Thomas TL, Hardin SH and Hall TC. 1995. Complete sequence of the binary vector Bin19. *Plant Mol. Biol.* 27 : 405-409.

Guerineau F and Mullineaux P. 1993. Plant transformation and expression vectors. In *Plant Molecular Biology Labfax*. Croy RRD (ed). BIOS Scientific Publishers, Blackwell Scientific Publications, pp 121-147.

Hanahan D. 1983. Studies on transformation of *Escherichia coli* with plasmids. *J. Mol. Biol.* 166 :557.

Holsters M, Dewaele D, Depicker A, Messenf E, Van Montagu M and Schell J. 1978. Transfection and transformation of *Agrobacterium tumefaciens*. *Mol. Gen. Genet.* 136 :181-187.

Horsch RB, Fry JE, Hoffmann NL, Eiholtz D, Rogers SG and Fraley RT. 1985. A simple and general method for transferring genes into plants. *Science* 227 :1229-1231.

Ishida Y, Saito H, Ohta S, Hiei Y, Komari T and Kumashiro T. 1996. High efficiency transformation of maize (*Zea mays* L.) mediated by *Agrobacterium tumefaciens*. *Nature Biotechnology* 14 :745-750.

Jefferson RA, Burgess SM and Hirsh D. 1986. b-glucuronidase from *Escherichia coli* as a gene-fusion marker. *Proc. Natl. Acad. Sci. (USA)* 83 :8447-8451.

Kay R, Chan A, Daly M and McPherson J. 1987. Duplication of CaMV 35S promoter sequences creates a strong enhancer for plant genes. *Science* 236 :1299-1302.

Klein TM, Wolf ED, Wu R and Sanford JC. 1987. High velocity microprojectiles for delivering nucleic acids into living cells. *Nature* 327 :70-73.

Komari T, Hiei Y, Yasuhito S, Murai N and Kumashiro T. 1996. Vectors carrying two separate T-DNAs for co-transformation of higher plants mediated by *Agrobacterium tumefaciens* and segregation of transformants free from selection markers. *Plant J.* 10 :165-174.

McElroy D, Blowers AD, Jenes B and Wu R. 1991. Construction of expression vectors based on the rice actin 1 (*Act1*) 5' region for the use in monocot transformation. *Mol. Gen. Genet.* 231 :150-160.

Michael SF. 1994. Mutagenesis by incorporation of a phosphorylated oligo for PCR amplification. *BioTechniques* 16 : 411-412.

Vain P, Yean H and Flament P. 1989. Enhancement of production and regeneration of embryogenic type II callus in *Zea mays* L. by AgNO_3 . *Plant Cell Tissue and Organ Culture* 18 :143-151.